

A Survey of Spherically Symmetric Spacetimes

Alan R. Parry*

October 22, 2012

Abstract

We present several different ways of describing a spherically symmetric spacetime and the resulting metrics. We then focus our discussion on an especially useful form of the metric of a spherically symmetric spacetime in polar-areal coordinates and its properties. In particular, we show how the metric component functions chosen are extremely compatible with notions in Newtonian mechanics. We also show the monotonicity of the Hawking mass in these coordinates. Finally, we discuss how these coordinates and the metric can be used to solve the spherically symmetric Einstein-Klein-Gordon equations and how this will be useful in our future work.

1 Introduction

Spherically symmetric spacetimes are an important case in the study of general relativity for a number of reasons. Foremost among them is that it is often a good starting point in the study of a problem in general relativity. For example, one of the first projects undergone in general relativity was to compute nontrivial spherically symmetric spacetimes that are exact solutions of the Einstein equation. This resulted in the discovery of the Schwarzschild spacetime, which is by far the most important spherically symmetric solution to date, and later Birkhoff's theorem about Ricci flat or vacuum spherically symmetric spacetimes.[14] Spherically symmetric spacetimes also create a situation where the dynamics of the system are less complicated by effectively reducing a 4-dimensional solution to a 2-dimensional one. This accessibility makes using spherically symmetric spacetimes all the more attractive as a starting point. Finally, while Birkhoff's theorem classifies all vacuum spherically symmetric spacetimes, there are still some nonvacuum spherically symmetric spacetimes that are interesting as well, both from a physical and mathematical standpoint, such as Dwarf spheroidal galaxies, which are dominated by their Dark Matter halos and also closely approximated by spherical symmetry[13], and spherically symmetric scalar fields.

Since spherically symmetric spacetimes are still of interest, it would be useful to find an efficient way to describe and model these spacetimes in a general sense. In particular, it would be useful to have a form of a general spherically symmetric metric that is well-suited to numerical evolutions of the Einstein equation. There are many possible forms of the metric of a general spherically symmetric spacetime to use and the purpose of this article is to first collect these

*Mathematics Department, Duke University, Box 90320, Durham, NC 27708, USA, alrparry@math.duke.edu

metrics, discuss the advantages and disadvantages of using each one, and then present a metric that is not only extremely well-suited to numerical evolutions, but also is described in terms that are very natural analogues to the low field or Newtonian limit. Most of these metrics can be described well within the established framework of numerical relativity and as such, we will use that framework in the following.

2 Metrics for a Spherically Symmetric Spacetime

The study of numerical relativity is devoted to devising ways of evolving the Einstein equation in order to solve for the components of the spacetime metric in different situations of interest as well as actually conducting such numerical experiments and comparing them to real data. This evolution usually takes place in a spacetime which is foliated by $t = \text{constant}$ spacelike hypersurfaces. The common method in numerical relativity is to decouple the time component from the space components into what is commonly called the (3+1)-formalism.[1, 2, 10]

The framework for this formalism is, as stated before, a spacetime, N , foliated by $t = \text{constant}$ spacelike hypersurfaces described by a Riemannian 3-metric γ which may change with time. Note that such a foliation is possible for any globally hyperbolic spacetime.[1, 14] Consider a coordinate chart on $U \subseteq N$, $\{t, x^1, x^2, x^3\}$ where the ∂_t is timelike and the ∂_{x^j} are all spacelike. Now consider an observer starting on the $t = t_0$ hypersurface at the coordinate $(t_0, \vec{x}_0^j)$. This observer then travels to another infinitesimally close hypersurface $t = t_0 + dt$ to the coordinate $(t_0 + dt, \vec{x}_0^j + d\vec{x})$ as in Figure 1 .




Figure 1: Infinitesimal distance in a $t = \text{constant}$ hypersurface foliated spacetime.

The observer has now traveled an infinitesimal distance ds . We can measure the square of this infinitesimal distance, ds^2 , using the analogue of Pythagorean's theorem and this will give us the line element form of the metric. If another observer travels normal to the hypersurface from $(t_0, \vec{x}_0^j)$ to the hypersurface $t = t_0 + dt$, since the normal direction is not necessarily the same direction as the t coordinate direction, it will arrive at the coordinate $(t_0 + dt, \vec{x}_0^j - \beta^j dt)$. We call the 3-vector field, β , the shift vector because it measures the spacelike shift of the coordinates while traveling normally. Note that the components of β can vary with all the coordinates. This normal observer, having traveled in a timelike direction, has experienced some proper time $d\tau$

which is some multiple of the change in time coordinate, that is,

$$d\tau = \alpha dt. \quad (1)$$

Hence the length of its normal movement from one surface to the other is αdt . The value of α can vary with all of the coordinates making it a function on the manifold, which is called the lapse function since it measures the lapse in proper time compared to coordinate time. To get the length in the spatial direction between where the normal observer ended up and where the original observer did, we need only to use the metric on the hypersurfaces and the difference of the two space coordinates. This difference, for each i , takes the form

$$(x_0^j + dx^j) - (x_0^j - \beta^j dt) = dx^j + \beta^j dt \quad (2)$$

and so the length squared of the spatial movement will be

$$\gamma_{jk}(dx^j + \beta^j dt)(dx^k + \beta^k dt) \quad (3)$$

where we have implemented the Einstein summation convention. Then using the generalized version of Pythagorean's theorem and recalling that t is a timelike direction we get that

$$ds^2 = -\alpha^2 dt^2 + \gamma_{jk}(dx^j + \beta^j dt)(dx^k + \beta^k dt), \quad (4)$$

which is the line element of the metric. This is the most general form of the metric for any foliated spacetime, that is, all metrics of a foliated spacetime can be written in this form. Note that we will often write g instead of ds^2 to refer interchangeably to the metric and the line element.

From this metric, and a choice of slicing condition, a complete system of partial differential equations that evolve the Einstein equation can be constructed. This system is often referred to as the ADM formulation of general relativity in reference to the authors of the paper in which it was first introduced.[4] It involves evolution equations of both the metric and the extrinsic curvature of the $t = \text{constant}$ hypersurfaces.[4, 6, 10] These equations are very commonly used in numerical relativity, but we find they overcomplicate the situation in spherical symmetry, which is why we have elected not to use this formulation of general relativity directly.

If we know more about the spacetime in question, we will be able to determine more of the components of the metric. In a spherically symmetric spacetime, with coordinates, r, θ, φ , chosen so that θ and φ are the polar-angular coordinates on the hypersurface, the shift vector must be completely radial so that the metric remains invariant under rotations. In this case, we will denote the radial component of the shift vector by simply β . Moreover, the 3-metric can be written as

$$\gamma = \gamma_{rr} dr^2 + \gamma_{\theta\theta} d\sigma^2. \quad (5)$$

where $d\sigma^2 = d\theta^2 + \sin^2 \theta d\varphi^2$ is the standard metric on the unit sphere. This implies that the most general spherically symmetric metric can be written in the form

$$\begin{aligned} g &= -\alpha^2 dt^2 + \gamma_{rr}(dr + \beta dt)(dr + \beta dt) + \gamma_{\theta\theta} d\sigma^2 \\ &= -(\alpha^2 - \gamma_{rr}\beta^2) dt^2 + \gamma_{rr}\beta(dr dt + dt dr) + \gamma_{rr} dr^2 + \gamma_{\theta\theta} d\sigma^2. \end{aligned} \quad (6)$$

Note that all the metric component functions can only depend on t and r due to spherical symmetry. For convenience, we will define two positive functions $a(t, r)$ and $q(t, r)$ so that

$$a(t, r)^2 = \gamma_{rr} \quad \text{and} \quad q(t, r)^2 = \gamma_{\theta\theta}. \quad (7)$$

Then we can rewrite (6) as

$$g = -(\alpha^2 - a^2\beta^2) dt^2 + a^2\beta(dr dt + dt dr) + a^2 dr^2 + q^2 d\sigma^2, \quad (8)$$

where α, a, β, q are functions of only t and r . It is important to note here that since we have not defined the coordinates t and r geometrically yet, there remain two degrees of freedom left in this metric. There are several different choices that can be made in this regard. We mention the most common here, but there is a very useful and more extensive list of several choices that can be made in a more general setting inourgoulhon's recent book.[1]

There are three rather common slicing conditions that are often used in many settings, all of which place a condition on the lapse function α . These conditions are the maximal slicing, harmonic slicing, and geodesic slicing conditions.

Under the maximal slicing condition, one requires that each $t = \text{constant}$ hypersurface be a maximal hypersurface, that is, it has zero mean curvature,

$$H = 0. \quad (9)$$

The metric stays of the form in equation (6), but since the evolution equations in the ADM formulation evolve the components of the second fundamental form and H is the trace of the second fundamental form, this places a constraint on some of the evolution variables, which can be used to simplify the evolution equations.[1, 5, 11] Note here that after making this choice, there remains one more degree of freedom which can be used to constrain the r coordinate.

Harmonic slicing requires that the coordinate function t be a harmonic function under the metric g . That is, $\square_g t = 0$, where $\square_g$ is the d'Alembertian or Laplacian operator with respect to the metric g . This is often accompanied with the condition that the hypersurfaces remain orthogonal to the time direction, which uses the remaining degree of freedom, and indeed some refer to both of these choices together as harmonic slicing. In the case that both conditions are satisfied, this would yield a metric of the form,

$$g = -\alpha^2 dt^2 + a^2 dr^2 + q^2 d\sigma^2 \quad (10)$$

with the added condition $\square_g t = 0$, which can be used to compute an evolution equation for the lapse function α . [7]

Geodesic slicing requires that movement along the curve $\xi = (t, 0, 0, 0)$, which is given in our coordinates, be geodesic, that is, coordinate observer worldlines are geodesics. This requirement is satisfied by choosing

$$\alpha = \text{constant} \quad \text{and} \quad \beta = 0. \quad (11)$$

However, the most reasonable choice for the constant is 1, since the condition $\alpha = 1$ on the lapse function has the added implication that normal observers proper time is the same as coordinate time and in fact, since $\beta = 0$, normal observers are coordinate observers.[1, 3, 5] This results in a metric of the form

$$g = -dt^2 + a^2 dr^2 + q^2 d\sigma^2. \quad (12)$$

While this choice seems very attractive at first, since it either eliminates or greatly simplifies the evolution equations, it does have a tendency to develop coordinate singularities when evolved in time and hence must be used with caution.[5] Note also that this requirement is sometimes reduced to simply $\alpha = 1$ without necessarily requiring the shift parameter or vector to vanish.

We can alternatively use the degrees of freedom to make choices concerning the r -coordinate. We will mention three such choices here that are standard in the study of spherically symmetric spacetimes.

The first is the choice of normal slicing. That is, choose the t coordinate so that the ∂_t vector field is always normal to the hypersurfaces. This choice makes all normal observers coordinate observers as well. This is equivalent to choosing the shift parameter or vector to be identically 0 and results in a metric of the form

$$g = -\alpha^2 dt^2 + a^2 dr^2 + q^2 d\sigma^2. \quad (13)$$

This choice has already been mentioned above as it is often coupled with harmonic or geodesic slicing conditions.[1] Since the coordinate vector fields on the hypersurfaces were already orthogonal, this results, as seen above, in a diagonal metric.

The next choice is to require that the metric on the hypersurfaces to be conformal to the flat metric. This amounts to choosing the function q in (8) to satisfy $q = ra$, which would make the metric become

$$g = -(\alpha^2 - a^2\beta^2) dt^2 + a^2\beta(dr dt + dt dr) + a^2(dr^2 + r^2 d\sigma^2). \quad (14)$$

This choice is referred to as isotropic coordinates. It is often coupled with the normal slicing choice above, which uses both of the degrees of freedom and results in a metric of the form

$$g = -\alpha^2 dt^2 + a^2(dr^2 + r^2 d\sigma^2). \quad (15)$$

A common example of the normal-isotropic case is the Schwarzschild metric in isotropic coordinates.[14]

The last choice we mention in the (3+1)-framework is to give the coordinate r geometric significance by choosing it to be the areal coordinate. That is, choose the coordinate r so that the area of each metric 2-sphere on the hypersurface is exactly $4\pi r^2$. This choice requires that $q = r$. Additionally, it is almost always accompanied with the normal slicing choice above, again using both degrees of freedom. This results in the polar-areal coordinates on a spherically symmetric spacetime and yields a metric of the form

$$g = -\alpha^2 dt^2 + a^2 dr^2 + r^2 d\sigma^2. \quad (16)$$

This chart is probably the most familiar form of a general spherically symmetric metric, and is the chart most commonly used when introducing the Schwarzschild spacetime. And rightly so, as it has some very clear advantages. For one, the r -coordinate's role is analogous to its role in a flat spacetime. Additionally, these coordinates give the Einstein curvature tensor a very simple form. However, they may not be well suited to dealing with high gravitational fields as we would likely run into the same limitations that the polar areal metric has in describing the Schwarzschild spacetime inside the Schwarzschild radius. This is not much of a problem for us, since we are mostly interested in the low field limit.

Finally, we present one more useful coordinate system and metric for a general spherically symmetric spacetime that does not fit into the (3+1)-formalism framework, nor does it depend on the ability to foliate the spacetime, but is useful from a theoretical standpoint nonetheless. In these coordinates, we still choose to use the polar-angular coordinates θ and φ to describe the rotations, but instead of separating the time and radial coordinates, we choose coordinates u and v such that ∂_u and ∂_v are nonparallel future pointing null vectors. This coordinate system is descriptively called null coordinates. In this case, the metric on the spacetime takes the form,

$$g = A(du dv + dv du) + Q^2 d\sigma^2 \quad (17)$$

where A and Q are functions of only u and v . Note that there are no extra degrees of freedom with this metric as all coordinates have been well-defined. While these coordinates are not very well suited to numerical evolutions, they can be very useful for theoretical discussions about the spherically symmetric spacetime. For example, in these coordinates, it is very straightforward to prove the monotonicity of the Hawking mass given the dominant energy condition. Additionally, it seems to perform well when in the presence of high gravitational fields. In the Schwarzschild case, these coordinates are known as the Kruskal coordinate system and is the system generally used to describe the region inside the Schwarzschild radius.[14]

All of these different coordinate choices and different forms of the metric on a spherically symmetric spacetime have advantages to them. However, for the problem of numerically evolving a spacetime metric in a low gravitational field, we find that the polar-areal coordinates are the best suited due to the very simple system one gets from the Einstein equation. As such, polar-areal coordinates is the coordinate system that we will use throughout this paper, but, in addition, we will introduce new variables that will give the metric a different form. This new form of the metric will result in the added advantages that the new metric functions have very clear analogues in the Newtonian or low-field limit and the Einstein curvature tensor will become even more simplified.

3 A Newtonian-Compatible Metric

Let N be a spherically symmetric, (3+1)-dimensional, asymptotically Schwarzschild (that is, it is Schwarzschild as $r \rightarrow \infty$) spacetime with a Lorentzian metric g and be foliated by $t = \text{constant}$ spacelike hypersurfaces. Choose the polar-areal coordinate system discussed above so that the t -coordinate direction is always normal to the hypersurfaces, the radial coordinate r is the areal coordinate (that is, so that the metric sphere of metric radius r always has a surface area of $4\pi r^2$), and θ and φ are the usual polar-angular coordinates on the hypersurface. In these coordinates, the metric g has the line element form

$$g = -\alpha(t, r)^2 dt^2 + a(t, r)^2 dr^2 + r^2 d\sigma^2 \quad (18)$$

Now we define the functions $V(t, r)$ and $M(t, r)$ as follows,

$$V = \log \alpha \quad M = \frac{r}{2} \left(\frac{a^2 - 1}{a^2} \right) \quad (19)$$

and rewrite the metric (18) in terms of these functions to obtain,

$$g = -e^{2V} dt^2 + \left(1 - \frac{2M}{r} \right)^{-1} dr^2 + r^2 d\sigma^2 \quad (20)$$

Note that we will always be interested in the low-field limit, where $M \ll r$, which necessarily requires that $M(t, 0) = 0$ for all t .

3.1 Compatibility with Newtonian Physics

Here, we wish to compute several important properties about this metric and the physical interpretation of both V and M . To physically interpret this metric, we will introduce the Einstein equation, but first, we present a property that doesn't require the Einstein equation, the proof of which can be found in Appendix A.

Proposition 3.1. *The function $M(t, r)$ is the spacetime Hawking mass of the metric sphere, $\Sigma_{t,r}$, for any given t and r .*

This suggests that we should interpret M as the mass of the system. However, there is an even stronger reason to do so, which we will investigate later. In order to prepare for that discussion, we need some preliminary results first about the relationship between the metric and its stress-energy tensor. To begin, consider this metric as a solution to the Einstein equation

$$G = 8\pi T. \quad (21)$$

To facilitate our discussion, we will compute the Einstein curvature tensor of this metric in certain directions. To that end, define the following unit vector fields.

$$\nu_t = e^{-V} \partial_t \quad \nu_r = \sqrt{1 - \frac{2M}{r}} \partial_r \quad \nu_\theta = \frac{1}{r} \partial_\theta \quad \nu_\varphi = \frac{1}{r \sin \theta} \partial_\varphi \quad (22)$$

Note that at every point, $p \in N$, except the coordinate singularities $r = 0$ and $\theta = \pm\pi$ (that is, all points where all the vector fields are well defined), these vector fields form an orthonormal basis of $T_p N$ and hence are a frame field. The Einstein curvature tensor is defined as

$$G = \text{Ric} - \frac{1}{2} Rg \quad (23)$$

where Ric and R are the Ricci curvature tensor and the scalar curvature of the spacetime respectively. Since both the Ricci curvature tensor and its trace R are present in this equation, we will need to know a few results about the Ricci curvature in these coordinates. We have the following lemma and subsequent corollary. While the proof of the corollary is short and is presented here, the proof of the lemma can be found in Appendix A.

Lemma 3.2. *The only nonzero components of the Ricci curvature tensor in the ν_η basis and the scalar curvature are as follows.*

1. $\text{Ric}(\nu_t, \nu_t) = \left(V_{rr} + V_r^2 + \frac{2V_r}{r} \right) \left(1 - \frac{2M}{r} \right) + \frac{V_r}{r} \left(\frac{M}{r} - M_r \right) + \frac{V_t M_t - M_{tt}}{r e^{2V}} \left(1 - \frac{2M}{r} \right)^{-1} - \frac{3M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r} \right)^{-2}$
2. $\text{Ric}(\nu_t, \nu_r) = \frac{2M_t}{r^2 e^V} \left(1 - \frac{2M}{r} \right)^{-1/2}$

$$\begin{aligned}
 3. \text{ Ric}(\nu_r, \nu_r) &= -(V_{rr} + V_r^2) \left(1 - \frac{2M}{r}\right) - \left(\frac{2}{r^2} + \frac{V_r}{r}\right) \left(\frac{M}{r} - M_r\right) \\
 &\quad - \frac{V_t M_t - M_{tt}}{r e^{2V}} \left(1 - \frac{2M}{r}\right)^{-1} + \frac{3M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r}\right)^{-2} \\
 4. \text{ Ric}(\nu_\theta, \nu_\theta) &= \text{ Ric}(\nu_\varphi, \nu_\varphi) = \frac{1}{r^2} \left(\frac{M}{r} + M_r\right) - \frac{V_r}{r} \left(1 - \frac{2M}{r}\right) \\
 5. R &= -2 \left(V_{rr} + V_r^2 + \frac{2V_r}{r}\right) \left(1 - \frac{2M}{r}\right) - \frac{2V_r}{r} \left(\frac{M}{r} - M_r\right) + \frac{4M_r}{r^2} \\
 &\quad - \frac{2(V_t M_t - M_{tt})}{r e^{2V}} \left(1 - \frac{2M}{r}\right)^{-1} + \frac{6M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r}\right)^{-2}
 \end{aligned}$$

Corollary 3.3. *The only nonzero components of the Einstein curvature tensor in the ν_η basis are as follows.*

$$\begin{aligned}
 1. G(\nu_t, \nu_t) &= \frac{2M_r}{r^2} \\
 2. G(\nu_t, \nu_r) &= \frac{2M_t}{r^2 e^V} \left(1 - \frac{2M}{r}\right)^{-1/2} \\
 3. G(\nu_r, \nu_r) &= -\frac{2M}{r^3} + \frac{2V_r}{r} \left(1 - \frac{2M}{r}\right) \\
 4. G(\nu_\theta, \nu_\theta) &= G(\nu_\varphi, \nu_\varphi) = \left(V_{rr} + V_r^2 + \frac{V_r}{r}\right) \left(1 - \frac{2M}{r}\right) + \left(\frac{M}{r} - M_r\right) \left(\frac{1}{r^2} + \frac{V_r}{r}\right) \\
 &\quad + \frac{V_t M_t - M_{tt}}{r e^{2V}} \left(1 - \frac{2M}{r}\right)^{-1} - \frac{3M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r}\right)^{-2}
 \end{aligned}$$

Proof. Since $\{\nu_t, \nu_r, \nu_\theta, \nu_\varphi\}$ form an orthonormal basis everywhere. We have the following.

$$G(\nu_t, \nu_t) = \text{Ric}(\nu_t, \nu_t) - \frac{R}{2} g(\nu_t, \nu_t) = \text{Ric}(\nu_t, \nu_t) + \frac{R}{2} \quad (24)$$

$$G(\nu_t, \nu_r) = \text{Ric}(\nu_t, \nu_r) - \frac{R}{2} g(\nu_t, \nu_r) = \text{Ric}(\nu_t, \nu_r) \quad (25)$$

$$G(\nu_r, \nu_r) = \text{Ric}(\nu_r, \nu_r) - \frac{R}{2} g(\nu_r, \nu_r) = \text{Ric}(\nu_r, \nu_r) - \frac{R}{2} \quad (26)$$

$$G(\nu_\theta, \nu_\theta) = \text{Ric}(\nu_\theta, \nu_\theta) - \frac{R}{2} g(\nu_\theta, \nu_\theta) = \text{Ric}(\nu_\theta, \nu_\theta) - \frac{R}{2} \quad (27)$$

$$\begin{aligned}
 G(\nu_\varphi, \nu_\varphi) &= \text{Ric}(\nu_\varphi, \nu_\varphi) - \frac{R}{2} g(\nu_\varphi, \nu_\varphi) = \text{Ric}(\nu_\varphi, \nu_\varphi) - \frac{R}{2} \\
 &= \text{Ric}(\nu_\theta, \nu_\theta) - \frac{R}{2} = G(\nu_\theta, \nu_\theta)
 \end{aligned} \quad (28)$$

We then use Lemma 3.2 to substitute in the values for Ric and R . The rest is algebra. These are the only nonzero components by Lemma 3.2, equation (23), and the fact that the ν_η basis is orthonormal. $\square$

Next, we define the function $\mu(t, r)$ to be the energy density of an observer at $p = (t, r)$ moving through the slices with 4-velocity ν_t . That is,

$$\mu = T(\nu_t, \nu_t). \quad (29)$$

We now have enough information to prove the following proposition, which contains the promised stronger reason for interpreting M as the mass inside each metric sphere.

Proposition 3.4. *For a fixed t and r , $M(t, r)$ is the flat integral of the energy density, $\mu(t, r)$, over the ball $E_{t,r}$ of radius r at time t .*

Proof. By the Einstein equation (21), equation (29), and Corollary 3.3, we have that

$$\begin{aligned} G(\nu_t, \nu_t) &= 8\pi T(\nu_t, \nu_t) \\ \frac{2M_r}{r^2} &= 8\pi\mu \\ M_r &= 4\pi r^2 \mu \end{aligned} \quad (30)$$

Since $\int_{\Sigma_{t,r}} dA = 4\pi r^2$, where $\Sigma_{t,r}$ is the sphere of radius r at time t , we have then that for a fixed t and r ,

$$M(t, r) = \int_0^r 4\pi s^2 \mu(t, s) ds = \int_0^r \int_{\Sigma_{t,s}} \mu(t, s) dA ds \quad (31)$$

$$= \int_0^r \int_0^{2\pi} \int_0^\pi \mu(t, s) s^2 \sin \theta d\theta d\varphi ds \quad (32)$$

$$= \int_{E_{t,r}} \mu(t, s) dV_0, \quad (33)$$

where $dV_0 = s^2 \sin \theta d\theta d\varphi ds$ is the flat volume form on the ball of radius s , and we have introduced s as a dummy integrating variable in the “ r ” position. Thus $M(t, r)$ is the flat volume integral of the energy density over the ball of radius r . $\square$

Note that M is not the integral of rhw energy density with respect to the metric’s volume form, $dV = (1 - 2M/s)^{-1/2} s^2 \sin \theta ds d\theta d\varphi$, but rather the following is true.

$$M(t, r) = \int_{E_{t,r}} \mu(t, s) \sqrt{1 - \frac{2M(t, s)}{s}} dV. \quad (34)$$

However, in the Newtonian limit, $M \ll r$, the above integral becomes approximately the integral of the energy density with respect to the metric’s volume form over the ball $E_{t,r}$ of radius r . Thus referring to $M(t, r)$ as the mass inside the metric sphere of radius r at time t not only makes sense from a geometrical point of view given the Hawking mass, but also from a physical point of view.

Next we define the function P to be the pressure in the stress-energy tensor for an observer at (t, r) , that is, let

$$P = T(\nu_r, \nu_r). \quad (35)$$

Then we can use Corollary 3.3 and the Einstein equation (21) to prove the following proposition.

Proposition 3.5. *In the Newtonian limit, where $P = 0$ and $M \ll r$, we have that $\Delta V = 4\pi\mu$, where Δ is the flat Laplacian on $\mathbb{R}^3$.*

Proof. By the Einstein equation (21), equation (35), and Corollary 3.3, we have that

$$\begin{aligned} G(\nu_r, \nu_r) &= 8\pi T(\nu_r, \nu_r) \\ -\frac{2M}{r^3} + \frac{2V_r}{r} \left(1 - \frac{2M}{r}\right) &= 8\pi P \\ \frac{2V_r}{r} &= \left(1 - \frac{2M}{r}\right)^{-1} \left(\frac{2M}{r^3} + 8\pi P\right) \\ r^2 V_r &= \left(1 - \frac{2M}{r}\right)^{-1} (M + 4\pi r^3 P). \end{aligned} \tag{36}$$

Note that in the Newtonian limit, where $P = 0$ and $M \ll r$, this equation is approximated by

$$r^2 V_r = M. \tag{37}$$

Also, the metric on the hypersurface, under these assumptions, is approximately the polar-area metric on $\mathbb{R}^3$. Note that the Laplacian on $\mathbb{R}^3$ of a spherically symmetric function f is given by

$$\Delta f = \frac{\partial^2 f}{\partial r^2} + \frac{2}{r} \frac{\partial f}{\partial r} = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial f}{\partial r} \right) \tag{38}$$

Then applying the operator $\frac{1}{r^2} \frac{\partial}{\partial r}$ to (37) and using equation (30) yields

$$\begin{aligned} \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 V_r) &= \frac{1}{r^2} \frac{\partial}{\partial r} (M) \\ \Delta V &= \frac{1}{r^2} M_r \\ \Delta V &= 4\pi\mu \end{aligned} \tag{39}$$

□

Equation (39) is Poisson's equation and is the defining equation of the gravitational potential in Newtonian mechanics. Moreover, equation (37) reduces to $V_r = M/r^2$ which yields the inverse square law for Newtonian gravity. Then we can interpret V as the analogue in our scenario of the Newtonian potential. The interpretation of M and V via Propositions 3.1, 3.4, and 3.5 are what we mean by saying that the metric (20) is Newtonian compatible.

3.2 Other Useful Properties of this Metric

In this section, we produce two additional useful results which are readily obtainable with these coordinates and metric functions. The first is the well-known result of the monotonicity of the Hawking mass in spherical symmetry, which is made particularly straightforward using this form of the metric. It follows almost immediately from Corollary 3.3.

Proposition 3.6. *If the spacetime satisfies the dominant energy condition that $G(X, Y) \geq 0$ for every future-pointing causal vector fields, X and Y , then, whenever $2M(t, r) \leq r$, the Hawking Mass, $M(t, r)$, is monotonically increasing in any non-timelike direction for which the radial coordinate increases.*

Proof. The vector field ν_t is the future-pointing timelike unit vector field in the t direction and the vector fields $\nu_r + \nu_t$ (future-pointing) and $\nu_r - \nu_t$ (past-pointing) are null vector fields in the null directions where the r coordinate increases. By Corollary 3.3, we have that

$$\begin{aligned}
 (\nu_r + \nu_t)(M) &= M_r \left(1 - \frac{2M}{r}\right)^{1/2} + \frac{M_t}{e^V} \\
 &= \frac{r^2}{2} \left(1 - \frac{2M}{r}\right)^{1/2} \left(\frac{2M_r}{r^2} + \frac{2M_t}{r^2 e^V} \left(1 - \frac{2M}{r}\right)^{-1/2} \right) \\
 &= \frac{r^2}{2} \left(1 - \frac{2M}{r}\right)^{1/2} (G(\nu_t, \nu_t) + G(\nu_t, \nu_r)) \\
 &= \frac{r^2}{2} \left(1 - \frac{2M}{r}\right)^{1/2} G(\nu_t, \nu_t + \nu_r)
 \end{aligned} \tag{40}$$

The last factor here is positive by the dominant energy condition since both ν_t and $\nu_t + \nu_r$ are future-pointing causal vector fields. Since $r^2 \geq 0$ and $2M \leq r$ everywhere, it must be that $(\nu_r + \nu_t)(M) \geq 0$ everywhere as well. By the same corollary, we also have

$$\begin{aligned}
 (\nu_r - \nu_t)(M) &= M_r \left(1 - \frac{2M}{r}\right)^{1/2} - \frac{M_t}{e^V} \\
 &= \frac{r^2}{2} \left(1 - \frac{2M}{r}\right)^{1/2} \left(\frac{2M_r}{r^2} - \frac{2M_t}{r^2 e^V} \left(1 - \frac{2M}{r}\right)^{-1/2} \right) \\
 &= \frac{r^2}{2} \left(1 - \frac{2M}{r}\right)^{1/2} (G(\nu_t, \nu_t) - G(\nu_t, \nu_r)) \\
 &= \frac{r^2}{2} \left(1 - \frac{2M}{r}\right)^{1/2} G(\nu_t, \nu_t - \nu_r)
 \end{aligned} \tag{41}$$

The last factor here is positive by the dominant energy condition since $\nu_t - \nu_r$ is also future-pointing causal. Then, as before, it must be that $(\nu_r - \nu_t)(M) \geq 0$ everywhere. Since both $(\nu_r + \nu_t)(M)$ and $(\nu_r - \nu_t)(M)$ are both nonnegative everywhere, any positive linear combination of the two is also nonnegative, which is the desired result. $\square$

This next property will be useful later when we consider the Einstein-Klein-Gordon system. By Corollary 3.3 and the Einstein equation, any stress-energy tensor T for this spacetime could have no other nonzero components than the corresponding components of the Einstein curvature tensor presented in Corollary 3.3. Thus we define the following functions, two of which have already been introduced in equations (29) and (35),

$$\mu(t, r) = T(\nu_t, \nu_t) \qquad \rho(t, r) = T(\nu_t, \nu_r) \tag{42a}$$

$$P(t, r) = T(\nu_r, \nu_r) \qquad Q(t, r) = T(\nu_\theta, \nu_\theta) = T(\nu_\varphi, \nu_\varphi) \tag{42b}$$

and these functions account for all the nonzero components of T in terms of the orthonormal frame $\{\nu_t, \nu_r, \nu_\theta, \nu_\varphi\}$. Then we have the following result which follows directly from the fact that $\nabla_g \cdot T = 0$, as with all stress-energy tensors, but whose lengthy proof we will leave to the appendix.

Proposition 3.7. *If the components of the stress-energy tensor T are known for all t and r and the functions $M(t, 0)$ and $V(t, 0)$ are known for all t , then solving the resulting initial value problem on the ODE's obtained from (30) and (36), namely*

$$M_r = 4\pi r^2 \mu \quad (43)$$

$$V_r = \left(1 - \frac{2M}{r}\right)^{-1} \left(\frac{M}{r^2} + 4\pi r P\right), \quad (44)$$

for each value of t , will solve the entire Einstein equation.

4 The Einstein-Klein-Gordon Equations

A good example of the usefulness of this choice of metric is in the evolution of the Einstein-Klein-Gordon equations, which is the Einstein equation in the presence of a nontrivial scalar field. In this paper, we will work specifically with a complex valued scalar field.

Consider a spherically symmetric spacetime N with Lorentzian metric g as described in the previous section. Let $f : N \rightarrow \mathbb{C}$ be a spherically symmetric complex valued scalar field. Then the Einstein-Klein Gordon equations in this case are

$$G = 8\pi\mu_0 \left(\frac{df \otimes d\bar{f} + d\bar{f} \otimes df}{\Upsilon^2} - \left(\frac{|df|^2}{\Upsilon^2} + |f|^2 \right) g \right) \quad (45)$$

$$\square_g f = \Upsilon^2 f \quad (46)$$

where μ_0 is simply some constant that controls the scale of the system and is not to be confused with the energy density $\mu(t, 0)$ at the central value. The value of μ_0 is unimportant to the qualitative behavior of the solutions and can be absorbed entirely into f , if desired. By contrast, the parameter Υ is a fundamental constant to the equations upon which the qualitative behavior of the solutions depend. The real valued version of these equations, that is, where f is a real valued scalar field, is derived in a paper by Bray.[9] Equations (45) and (46) are the Euler-Lagrange equations of the action involving the scalar field given in Bray's paper if the scalar field is assumed to be complex valued instead of real valued (this follows the Lagrangian formulation of general relativity given in an appendix in Wald's book[14]). One can readily see that we recover the equations in Bray's paper, if we assume that f is real valued, which implies that $\bar{f} = f$. Note also that equations (21) and (45) imply that for a complex scalar field

$$T = \mu_0 \left(\frac{df \otimes d\bar{f} + d\bar{f} \otimes df}{\Upsilon^2} - \left(\frac{|df|^2}{\Upsilon^2} + |f|^2 \right) g \right). \quad (47)$$

With these equations, we will construct a system of partial differential equations which can be used to numerically evolve the scalar field f and the metric from consistent initial data. Since

in the previous section we have already computed the components of the Einstein curvature tensor, in order to construct such a system of PDEs, we need to compute the components of the stress energy tensor and the Laplacian in the metric g . Before we do, we will define the function, $p(t, r)$, by the equation

$$p = f_t e^{-V} \left(1 - \frac{2M}{r}\right)^{-1/2}. \quad (48)$$

This is done to make the resulting system of PDEs first order in time and results in a more convenient choice than choosing only $p = f_t$. We now have the following lemmas.

Lemma 4.1. *The following are true.*

1. $T(\nu_t, \nu_t) = \mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right)$
2. $T(\nu_t, \nu_r) = \frac{2\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \text{Re}(f_r \bar{p})$
3. $T(\nu_r, \nu_r) = \mu_0 \left(-|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right)$
4. $T(\nu_\theta, \nu_\theta) = T(\nu_\varphi, \nu_\varphi) = -\mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} \right)$

The proof of this lemma is left to the appendix.

Lemma 4.2. *The components of the Einstein equation in the ν_η directions result in the following PDEs.*

$$M_r = 4\pi r^2 \mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \quad (49)$$

$$M_t = \frac{8\pi r^2 \mu_0 e^V}{\Upsilon^2} \left(1 - \frac{2M}{r}\right)^{3/2} \text{Re}(f_r \bar{p}) \quad (50)$$

$$V_r = \left(1 - \frac{2M}{r}\right)^{-1} \left(\frac{M}{r^2} - 4\pi r \mu_0 \left(|f|^2 - \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \right) \quad (51)$$

$$\begin{aligned} V_t M_t = -r e^{2V} & \left[\left(V_{rr} + V_r^2 + \frac{V_r}{r} \right) \left(1 - \frac{2M}{r}\right)^2 + \left(\frac{M}{r} - M_r \right) \left(\frac{1}{r^2} + \frac{V_r}{r} \right) \left(1 - \frac{2M}{r}\right) \right. \\ & \left. - \frac{M_{tt}}{r e^{2V}} - \frac{3M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r}\right)^{-1} + 8\pi \mu_0 \left(1 - \frac{2M}{r}\right) \left(|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} \right) \right] \end{aligned} \quad (52)$$

Proof. This follows directly from the Einstein equation (21), Corollary 3.3, and Lemma 4.1 and the algebra necessary to solve for each of the quantities above. Equation (49) follows from the ν_t, ν_t component of the Einstein equation, (50) follows from the ν_t, ν_r component, (51) from the ν_r, ν_r component, and (52) from the ν_θ, ν_θ or ν_φ, ν_φ component since they're identical. $\square$

The Klein-Gordon equation and the fact that

$$\square_g f = \frac{1}{\sqrt{|g|}} \partial_\eta \left(\sqrt{|g|} g^{\eta\omega} \partial_\omega f \right). \quad (53)$$

yield the following lemma.

Lemma 4.3. *The Klein-Gordon equation in this metric yields the following PDE.*

$$p_t = e^V \left(-\Upsilon^2 f \left(1 - \frac{2M}{r} \right)^{-1/2} + \frac{2f_r}{r} \sqrt{1 - \frac{2M}{r}} \right) + \partial_r \left(e^V f_r \sqrt{1 - \frac{2M}{r}} \right) \quad (54)$$

Proof. To compute the Laplacian, we'll first need the quantity $\sqrt{|g|}$. We have

$$\sqrt{|g|} = \sqrt{e^{2V} \left(1 - \frac{2M}{r} \right)^{-1} r^4 \sin^2 \theta} = e^V \left(1 - \frac{2M}{r} \right)^{-1/2} r^2 \sin \theta \quad (55)$$

Then since $f = f(t, r)$ and g is diagonal, we have by equation (53)

$$\begin{aligned} \square_g f &= \frac{1}{e^V r^2 \sin \theta} \sqrt{1 - \frac{2M}{r}} \left(\partial_t \left(e^V \left(1 - \frac{2M}{r} \right)^{-1/2} r^2 \sin \theta g^{tt} f_t \right) \right. \\ &\quad \left. + \partial_r \left(e^V \left(1 - \frac{2M}{r} \right)^{-1/2} r^2 \sin \theta g^{rr} f_r \right) \right) \\ &= \frac{1}{e^V r^2 \sin \theta} \sqrt{1 - \frac{2M}{r}} \left(\partial_t \left(-e^{-V} \left(1 - \frac{2M}{r} \right)^{-1/2} r^2 \sin \theta f_t \right) \right. \\ &\quad \left. + \partial_r \left(e^V \sqrt{1 - \frac{2M}{r}} r^2 \sin \theta f_r \right) \right) \\ &= \frac{1}{e^V r^2 \sin \theta} \sqrt{1 - \frac{2M}{r}} \left(\partial_t (-pr^2 \sin \theta) \right. \\ &\quad \left. + 2re^V \sqrt{1 - \frac{2M}{r}} \sin \theta f_r + r^2 \sin \theta \partial_r \left(e^V f_r \sqrt{1 - \frac{2M}{r}} \right) \right) \\ &= -p_t e^{-V} \sqrt{1 - \frac{2M}{r}} + \frac{2f_r}{r} \left(1 - \frac{2M}{r} \right) + e^{-V} \sqrt{1 - \frac{2M}{r}} \partial_r \left(e^V f_r \sqrt{1 - \frac{2M}{r}} \right) \end{aligned} \quad (56)$$

Then the Klein-Gordon equation (46) becomes

$$-p_t e^{-V} \sqrt{1 - \frac{2M}{r}} + \frac{2f_r}{r} \left(1 - \frac{2M}{r} \right) + e^{-V} \sqrt{1 - \frac{2M}{r}} \partial_r \left(e^V f_r \sqrt{1 - \frac{2M}{r}} \right) = \Upsilon^2 f \quad (57)$$

Solving for p_t yields the desired result. $\square$

Then rewrite (48) as

$$f_t = pe^V \sqrt{1 - \frac{2M}{r}} \quad (58)$$

to yield an evolution equation for f .

We now include a lemma that shows that the T given in (47) has the property of all stress-energy tensors that $\nabla_g \cdot T = 0$.

Lemma 4.4. *Equation (46) implies that $\nabla_g \cdot T = 0$, where T is given by (47).*

This lemma's proof is also included in the appendix. Now we can prove the following.

Lemma 4.5. *If f, p, M, V solve (49), (51), (54), and (58), then they also solve (50) and (52).*

Proof. This follows from equation (42), Lemmas 4.1, 4.2, 4.4, Proposition 3.7, and the fact that we can use (54) and (58) to evolve the components of the stress-energy tensor through time as long as we use (49) and (51) to compute the metric components at each new time. If we do these two procedures simultaneously, it is effectively the same as knowing the components of the stress-energy tensor for all values of t , which is required by the hypothesis of Proposition 3.7. $\square$

Lemmas 4.2, 4.3, and 4.5 and equation (58) prove the following proposition.

Proposition 4.6. *Solving the following PDEs for $f(t, r)$, $p(t, r)$, $V(t, r)$, and $M(t, r)$ with consistent initial data is necessary and sufficient to solve the Einstein-Klein-Gordon equations (45) and (46) with a metric of the form (20) on the same initial data.*

$$M_r = 4\pi r^2 \mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \quad (59a)$$

$$V_r = \left(1 - \frac{2M}{r}\right)^{-1} \left(\frac{M}{r^2} - 4\pi r \mu_0 \left(|f|^2 - \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \right) \quad (59b)$$

$$f_t = pe^V \sqrt{1 - \frac{2M}{r}} \quad (59c)$$

$$p_t = e^V \left(-\Upsilon^2 f \left(1 - \frac{2M}{r}\right)^{-1/2} + \frac{2f_r}{r} \sqrt{1 - \frac{2M}{r}} \right) + \partial_r \left(e^V f_r \sqrt{1 - \frac{2M}{r}} \right) \quad (59d)$$

5 Future Work

Of all the different coordinate systems and metric functions that we have used, this Newtonian compatible choice yields by far the simplest system of PDEs that we have come across that solve the spherically symmetric Einstein-Klein-Gordon equations. Solutions to the Einstein-Klein-Gordon equations could possibly yield a geometric description of dark matter. This form of dark matter, which we call wave dark matter, has been studied for years, but has recently been well motivated geometrically by Bray.[9] In much of our future work, we plan to study the geometric and gravitational properties of solutions to the Einstein-Klein-Gordon equations in order to compare them to observations.

One of the best places to start is with dwarf spheroidal galaxies, which are small, roughly spherically symmetric, and dark matter dominated. The system (59) which solves the spherically symmetric Einstein-Klein-Gordon equations will be particularly useful in this regard. We will use this system to study both static and non-static solutions to the Einstein-Klein-Gordon equations and compare their predictions to observations of the velocities of the stars in these galaxies. To this end, we have created Matlab code to take initial conditions of the scalar field $f(0, r) = f_0(r)$ and $p(0, r) = p_0(r)$ and evolve the entire Einstein-Klein-Gordon system indefinitely. The code works on relatively simple finite difference techniques and will be discussed in another work. The results of these endeavors will help begin to answer the question of whether or not wave dark matter correctly describes the gravitational effects of dark matter and will also help us determine the value of Υ , the fundamental constant in the Einstein-Klein-Gordon equations.

6 Acknowledgments

The author would like to thank Hubert Bray for several extremely useful discussions on this topic. We also gratefully acknowledge the support of the National Science Foundation grant # DMS-1007063.

A Proofs

In this appendix, we present the proofs which were omitted in the above sections. Every proof of a proposition or lemma from above will refer back to the number of the statement it is proving. We also introduce a new lemma here which is useful for proving some of the above statements. We will present all of the omitted proofs here in the order with which their corresponding statements are given above.

Proof of Proposition 3.1. The Hawking mass of a metric sphere is defined as

$$m_H(\Sigma_{t,r}) = \sqrt{\frac{|\Sigma_{t,r}|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma_{t,r}} g(\vec{H}, \vec{H}) dA \right) \quad (60)$$

where $|\Sigma_{t,r}|$ is the surface area of the metric sphere, $\vec{H}$ is the mean curvature vector of the sphere in the spacetime, and dA is the volume element on the sphere.[8, 12] It has already been established that $|\Sigma_{t,r}| = 4\pi r^2$, but it is also easily computed in this metric since

$$dA = \sqrt{|d\sigma^2|} d\theta d\varphi = \sqrt{r^4 \sin^2 \theta} d\theta d\varphi = r^2 \sin \theta d\theta d\varphi, \quad (61)$$

which yields that

$$|\Sigma_{t,r}| = \int_{\Sigma_{t,r}} dA = \int_0^{2\pi} \int_0^\pi r^2 \sin \theta d\theta d\varphi = \int_0^{2\pi} 2r^2 d\varphi = 4\pi r^2. \quad (62)$$

To compute the mean curvature vector, $\vec{H}$, we have that

$$\vec{H} = \gamma^{jk} \Pi(\partial_j, \partial_k) \quad (63)$$

where $j, k \in \{\theta, \varphi\}$, γ is the metric on $\Sigma_{t,r}$ (note that we have reused the variable γ here to refer to the metric on the sphere and not to the metric on the $t = \text{constant}$ hypersurface as we did in the section describing the framework of numerical relativity), and Π is the second fundamental form tensor, which sends a pair of vectors tangent to the sphere to a vector normal to the sphere. It is defined as follows for all $X, Y \in T\Sigma_{t,r}$

$$\Pi(X, Y) = \nabla_X Y - {}^2\nabla_X Y \quad (64)$$

where ∇ is the covariant derivative operator on N and ${}^2\nabla$ is the induced covariant derivative operator on $\Sigma_{t,r}$. Since γ is diagonal, we only need to concern ourselves with the diagonal components of this tensor in order to compute $\vec{H}$. To perform these computations, first note that the Christoffel symbols on the sphere in these coordinates have the following property (note that a superscript of 2 will always denote that that quantity corresponds to the sphere, also for the following computations, roman letters will denote subscripts in $\{\theta, \varphi\}$, while Greek letters will denote subscripts ranging over all four coordinates).

$$\Gamma_{jk}^\ell = \frac{1}{2}g^{\ell\eta}(g_{j\eta,k} + g_{k\eta,j} - g_{jk,\eta}) = \frac{1}{2}g^{\ell m}(g_{jm,k} + g_{km,j} - g_{jk,m}) = \frac{1}{2}\gamma^{\ell m}(g_{jm,k} + g_{km,j} - g_{jk,m}) = {}^2\Gamma_{jk}^\ell \quad (65)$$

since both the radial and time directions are normal to the sphere. So we need not distinguish between the Christoffel symbols for the metric on the sphere and those for the entire manifold. We have then that

$$\Pi(\partial_\theta, \partial_\theta) = \nabla_\theta \partial_\theta - {}^2\nabla_\theta \partial_\theta = \Gamma_{\theta\theta}^\eta \partial_\eta - {}^2\Gamma_{\theta\theta}^k \partial_k = \Gamma_{\theta\theta}^t \partial_t + \Gamma_{\theta\theta}^r \partial_r \quad (66)$$

and similarly

$$\Pi(\partial_\varphi, \partial_\varphi) = \nabla_\varphi \partial_\varphi - {}^2\nabla_\varphi \partial_\varphi = \Gamma_{\varphi\varphi}^\eta \partial_\eta - {}^2\Gamma_{\varphi\varphi}^k \partial_k = \Gamma_{\varphi\varphi}^t \partial_t + \Gamma_{\varphi\varphi}^r \partial_r \quad (67)$$

Then we need to compute the above Christoffel symbols. We have that

$$\Gamma_{\theta\theta}^t = \frac{1}{2}g^{t\eta}(g_{\theta\eta,\theta} + g_{\theta\eta,\theta} - g_{\theta\theta,\eta}) = \frac{1}{2}g^{tt}(-g_{\theta\theta,t}) = \frac{1}{2e^{2V}}(-\partial_t(r^2)) = 0 \quad (68a)$$

$$\begin{aligned} \Gamma_{\theta\theta}^r &= \frac{1}{2}g^{r\eta}(g_{\theta\eta,\theta} + g_{\theta\eta,\theta} - g_{\theta\theta,\eta}) = \frac{1}{2}g^{rr}(-g_{\theta\theta,r}) \\ &= \frac{1}{2}\left(1 - \frac{2M}{r}\right)(-\partial_r(r^2)) = -r\left(1 - \frac{2M}{r}\right) \end{aligned} \quad (68b)$$

$$\Gamma_{\varphi\varphi}^t = \frac{1}{2}g^{t\eta}(g_{\varphi\eta,\varphi} + g_{\varphi\eta,\varphi} - g_{\varphi\varphi,\eta}) = \frac{1}{2}g^{tt}(-g_{\varphi\varphi,t}) = \frac{1}{2e^{2V}}(-\partial_t(r^2 \sin^2 \theta)) = 0 \quad (68c)$$

$$\begin{aligned} \Gamma_{\varphi\varphi}^r &= \frac{1}{2}g^{r\eta}(g_{\varphi\eta,\varphi} + g_{\varphi\eta,\varphi} - g_{\varphi\varphi,\eta}) = \frac{1}{2}g^{rr}(-g_{\varphi\varphi,r}) \\ &= \frac{1}{2}\left(1 - \frac{2M}{r}\right)(-\partial_r(r^2 \sin^2 \theta)) = -r\left(1 - \frac{2M}{r}\right)\sin^2 \theta. \end{aligned} \quad (68d)$$

Then we have that

$$\Pi(\partial_\theta, \partial_\theta) = \Gamma_{\theta\theta}^t \partial_t + \Gamma_{\theta\theta}^r \partial_r = -r\left(1 - \frac{2M}{r}\right) \partial_r \quad (69)$$

$$\Pi(\partial_\varphi, \partial_\varphi) = \Gamma_{\varphi\varphi}^t \partial_t + \Gamma_{\varphi\varphi}^r \partial_r = -r\left(1 - \frac{2M}{r}\right) \sin^2 \theta \partial_r \quad (70)$$

This makes (63) become

$$\begin{aligned}
 \vec{H} &= \gamma^{\theta\theta} \Pi(\partial_\theta, \partial_\theta) + g^{\varphi\varphi} \Pi(\partial_\varphi, \partial_\varphi) \\
 &= -\frac{1}{r^2} \cdot r \left(1 - \frac{2M}{r}\right) \partial_r - \frac{1}{r^2 \sin^2 \theta} \cdot r \left(1 - \frac{2M}{r}\right) \sin^2 \theta \partial_r \\
 &= -\frac{2}{r} \left(1 - \frac{2M}{r}\right) \partial_r
 \end{aligned} \tag{71}$$

Finally, we can compute the Hawking mass as follows.

$$\begin{aligned}
 m_H(\Sigma_{t,r}) &= \sqrt{\frac{|\Sigma_{t,r}|}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma_{t,r}} g(\vec{H}, \vec{H}) dA\right) \\
 &= \sqrt{\frac{4\pi r^2}{16\pi}} \left(1 - \frac{1}{16\pi} \int_{\Sigma_{t,r}} g\left(-\frac{2}{r} \left(1 - \frac{2M}{r}\right) \partial_r, -\frac{2}{r} \left(1 - \frac{2M}{r}\right) \partial_r\right) dA\right) \\
 &= \frac{r}{2} \left(1 - \frac{1}{16\pi} \int_0^{2\pi} \int_0^\pi \frac{4}{r^2} \left(1 - \frac{2M}{r}\right)^2 g(\partial_r, \partial_r) r^2 \sin \theta d\theta d\varphi\right) \\
 &= \frac{r}{2} \left(1 - \frac{1}{16\pi} \int_0^{2\pi} \int_0^\pi 4 \left(1 - \frac{2M}{r}\right) \sin \theta d\theta d\varphi\right) \\
 &= \frac{r}{2} \left(1 - \frac{1}{4\pi} \left(1 - \frac{2M}{r}\right) \int_0^{2\pi} \int_0^\pi \sin \theta d\theta d\varphi\right) \\
 &= \frac{r}{2} \left(1 - \frac{1}{4\pi} \left(1 - \frac{2M}{r}\right) (4\pi)\right) \\
 &= \frac{r}{2} \left(\frac{2M}{r}\right) \\
 m_H(\Sigma_{t,r}) &= M
 \end{aligned} \tag{72}$$

□

To prove Lemma 3.2, we will utilize the following additional lemma.

Lemma A.1. *For all $\eta \in \{t, r, \theta, \varphi\}$, let ν_η be as defined by equation (22). Then the following are true.*

1. *For all $\eta \in \{t, r, \theta, \varphi\}$, $\text{Ric}(\nu_\eta, \nu_\eta) = g(\nu_\eta, \nu_\eta) g^{\eta\eta} \text{Ric}_{\eta\eta}$.*
2. $R = \sum_{\eta} g(\nu_\eta, \nu_\eta) \text{Ric}(\nu_\eta, \nu_\eta)$

Proof. Since $\{\nu_\eta\}$ is a frame field, we have that

$$\text{Ric}(\nu_t, \nu_t) = e^{-2V} \text{Ric}(\partial_t, \partial_t) = -g^{tt} \text{Ric}_{tt} = g(\nu_t, \nu_t) g^{tt} \text{Ric}_{tt}. \tag{73}$$

By similar arguments, we have that for $\eta \in \{t, r, \theta, \varphi\}$,

$$\text{Ric}(\nu_\eta, \nu_\eta) = g(\nu_\eta, \nu_\eta) g^{\eta\eta} \text{Ric}_{\eta\eta} \tag{74}$$

which proves 1. To prove 2, we use the first result and find that since g is diagonal and $g(\nu_\eta, \nu_\eta) = \pm 1$, we have that

$$R = \sum_{\eta\omega} g^{\eta\omega} \text{Ric}_{\eta\omega} = \sum_{\eta} g^{\eta\eta} \text{Ric}_{\eta\eta} = \sum_{\eta} \frac{\text{Ric}(\nu_\eta, \nu_\eta)}{g(\nu_\eta, \nu_\eta)} = \sum_{\eta} g(\nu_\eta, \nu_\eta) \text{Ric}(\nu_\eta, \nu_\eta). \quad (75)$$

Fact 2 also follows from a well understood more general property of frame fields and traces of tensors. $\square$

With this, we prove Lemma 3.2.

Proof of Lemma 3.2. In this proof, all indices range over the coordinates $\{t, r, \theta, \varphi\}$. By Lemma A.1 and the well known formula for the components of the Riemann curvature tensor, $\mathbf{R}$, in terms of the Christoffel symbols

$$\mathbf{R}_{jkl}{}^m = \Gamma_{jl}{}^m{}_{,k} - \Gamma_{kl}{}^m{}_{,j} + \Gamma_{jl}{}^s \Gamma_{ks}{}^m - \Gamma_{kl}{}^s \Gamma_{js}{}^m. \quad (76)$$

Since $\text{Ric}_{jk} = \mathbf{R}_{j\ell k}{}^\ell$, we have that

$$\begin{aligned} \text{Ric}(\nu_t, \nu_t) &= g(\nu_t, \nu_t) g^{tt} \text{Ric}_{tt} \\ &= e^{-2V} (\Gamma_{tt}{}^k{}_{,k} - \Gamma_{kt}{}^k{}_{,t} + \Gamma_{tt}{}^s \Gamma_{ks}{}^k - \Gamma_{kt}{}^s \Gamma_{ts}{}^k) \end{aligned} \quad (77a)$$

$$\begin{aligned} \text{Ric}(\nu_t, \nu_r) &= e^{-V} \sqrt{1 - \frac{2M}{r}} \text{Ric}_{tr} \\ &= e^{-V} \sqrt{1 - \frac{2M}{r}} (\Gamma_{tr}{}^k{}_{,k} - \Gamma_{kr}{}^k{}_{,t} + \Gamma_{tr}{}^s \Gamma_{ks}{}^k - \Gamma_{kr}{}^s \Gamma_{ts}{}^k) \end{aligned} \quad (77b)$$

$$\begin{aligned} \text{Ric}(\nu_r, \nu_r) &= g(\nu_r, \nu_r) g^{rr} \text{Ric}_{rr} = \left(1 - \frac{2M}{r}\right) \text{Ric}_{rr} \\ &= \left(1 - \frac{2M}{r}\right) (\Gamma_{rr}{}^k{}_{,k} - \Gamma_{kr}{}^k{}_{,r} + \Gamma_{rr}{}^s \Gamma_{ks}{}^k - \Gamma_{kr}{}^s \Gamma_{rs}{}^k) \end{aligned} \quad (77c)$$

$$\begin{aligned} \text{Ric}(\nu_\theta, \nu_\theta) &= g(\nu_\theta, \nu_\theta) g^{\theta\theta} \text{Ric}_{\theta\theta} = \frac{1}{r^2} \text{Ric}_{\theta\theta} \\ &= \frac{1}{r^2} (\Gamma_{\theta\theta}{}^k{}_{,k} - \Gamma_{k\theta}{}^k{}_{,\theta} + \Gamma_{\theta\theta}{}^s \Gamma_{ks}{}^k - \Gamma_{k\theta}{}^s \Gamma_{\theta s}{}^k) \end{aligned} \quad (77d)$$

$$\begin{aligned} \text{Ric}(\nu_\varphi, \nu_\varphi) &= g(\nu_\varphi, \nu_\varphi) g^{\varphi\varphi} \text{Ric}_{\varphi\varphi} = \frac{1}{r^2 \sin^2 \theta} \text{Ric}_{\varphi\varphi} \\ &= \frac{1}{r^2 \sin^2 \theta} (\Gamma_{\varphi\varphi}{}^k{}_{,k} - \Gamma_{k\varphi}{}^k{}_{,\varphi} + \Gamma_{\varphi\varphi}{}^s \Gamma_{ks}{}^k - \Gamma_{k\varphi}{}^s \Gamma_{\varphi s}{}^k). \end{aligned} \quad (77e)$$

Then to compute these, we need the Christoffel symbols. For convenience, we will write the Christoffel symbols in four matrices, Γ^t , Γ^r , Γ^θ , and Γ^φ , which are the 4×4 matrices of Christoffel symbols corresponding to a given fixed upper index. To do so, recall, that the formula for the Christoffel symbols in terms of the metric components is the following

$$\Gamma_{jk}{}^\ell = \frac{1}{2} g^{\ell m} (g_{jm,k} + g_{km,j} - g_{jk,m}) \quad (78)$$

Using (78) and the fact that g is diagonal, we have that

$$\begin{aligned}\Gamma_{jk}^t &= \frac{1}{2}g^{tm}(g_{jm,k} + g_{km,j} - g_{jk,m}) = \frac{1}{2}g^{tt}(g_{jt,k} + g_{kt,j} - g_{jk,t}) \\ &= -\frac{1}{2e^{2V}}(g_{jt,k} + g_{kt,j} - g_{jk,t})\end{aligned}\quad (79a)$$

$$\begin{aligned}\Gamma_{jk}^r &= \frac{1}{2}g^{rm}(g_{jm,k} + g_{km,j} - g_{jk,m}) = \frac{1}{2}g^{rr}(g_{jr,k} + g_{kr,j} - g_{jk,r}) \\ &= \frac{1}{2}\left(1 - \frac{2M}{r}\right)(g_{jr,k} + g_{kr,j} - g_{jk,r})\end{aligned}\quad (79b)$$

$$\begin{aligned}\Gamma_{jk}^\theta &= \frac{1}{2}g^{\theta m}(g_{jm,k} + g_{km,j} - g_{jk,m}) = \frac{1}{2}g^{\theta\theta}(g_{j\theta,k} + g_{k\theta,j} - g_{jk,\theta}) \\ &= \frac{1}{2r^2}(g_{j\theta,k} + g_{k\theta,j} - g_{jk,\theta})\end{aligned}\quad (79c)$$

$$\begin{aligned}\Gamma_{jk}^\varphi &= \frac{1}{2}g^{\varphi m}(g_{jm,k} + g_{km,j} - g_{jk,m}) = \frac{1}{2}g^{\varphi\varphi}(g_{j\varphi,k} + g_{k\varphi,j} - g_{jk,\varphi}) \\ &= \frac{1}{2r^2 \sin^2 \theta}(g_{j\varphi,k} + g_{k\varphi,j} - g_{jk,\varphi}) = \frac{1}{2r^2 \sin^2 \theta}(g_{j\varphi,k} + g_{k\varphi,j}).\end{aligned}\quad (79d)$$

The last line is due to the fact that none of the metric components depend on φ . To help compute these quantities, it would be useful to note the following.

$$\begin{aligned}g_{tt,t} &= \partial_t(-e^{2V}) = -2V_t e^{2V} & g_{tt,\theta} &= \partial_\theta(-e^{2V}) = 0 \\ g_{tt,r} &= \partial_r(-e^{2V}) = -2V_r e^{2V} & g_{tt,\varphi} &= \partial_\varphi(-e^{2V}) = 0\end{aligned}\quad (80a)$$

$$\begin{aligned}g_{rr,t} &= \partial_t\left(1 - \frac{2M}{r}\right)^{-1} & g_{rr,\theta} &= \partial_\theta\left(1 - \frac{2M}{r}\right)^{-1} = 0 \\ &= \left(1 - \frac{2M}{r}\right)^{-2}\left(\frac{2M_t}{r}\right) \\ g_{rr,r} &= \partial_r\left(1 - \frac{2M}{r}\right)^{-1} & g_{rr,\varphi} &= \partial_\varphi\left(1 - \frac{2M}{r}\right)^{-1} = 0 \\ &= \left(1 - \frac{2M}{r}\right)^{-2}\left(\frac{2M_r}{r} - \frac{2M}{r^2}\right)\end{aligned}\quad (80b)$$

$$\begin{aligned}g_{\theta\theta,t} &= \partial_t(r^2) = 0 & g_{\theta\theta,\theta} &= \partial_\theta(r^2) = 0 \\ g_{\theta\theta,r} &= \partial_r(r^2) = 2r & g_{\theta\theta,\varphi} &= \partial_\varphi(r^2) = 0\end{aligned}\quad (80c)$$

$$\begin{aligned}g_{\varphi\varphi,t} &= \partial_t(r^2 \sin^2 \theta) = 0 & g_{\varphi\varphi,\theta} &= \partial_\theta(r^2 \sin^2 \theta) = 2r^2 \sin \theta \cos \theta \\ g_{\varphi\varphi,r} &= \partial_r(r^2 \sin^2 \theta) = 2r \sin^2 \theta & g_{\varphi\varphi,\varphi} &= \partial_\varphi(r^2 \sin^2 \theta) = 0.\end{aligned}\quad (80d)$$

All the other metric components are 0. Using (79) and (80), it is a series of straightforward algebra computations to obtain the following.

$$\Gamma^t = \begin{pmatrix} V_t & V_r & 0 & 0 \\ V_r & \frac{M_t}{re^{2V}}\left(1 - \frac{2M}{r}\right)^{-2} & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}\quad (81a)$$

$$\Gamma^r = \begin{pmatrix} V_r e^{2V} \left(1 - \frac{2M}{r}\right) & \frac{M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} & 0 & 0 \\ \frac{M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} & \left(1 - \frac{2M}{r}\right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2}\right) & 0 & 0 \\ 0 & 0 & -r \left(1 - \frac{2M}{r}\right) & 0 \\ 0 & 0 & 0 & -r \sin^2 \theta \left(1 - \frac{2M}{r}\right) \end{pmatrix} \quad (81b)$$

$$\Gamma^\theta = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{r} & 0 \\ 0 & \frac{1}{r} & 0 & 0 \\ 0 & 0 & 0 & -\sin \theta \cos \theta \end{pmatrix} \quad (81c)$$

$$\Gamma^\varphi = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{r} \\ 0 & 0 & 0 & \frac{\cos \theta}{\sin \theta} \\ 0 & \frac{1}{r} & \frac{\cos \theta}{\sin \theta} & 0 \end{pmatrix} \quad (81d)$$

Using (81), we can now continue the computations started in (77).

$$\begin{aligned} \text{Ric}(\nu_t, \nu_t) &= e^{-2V} \left(\Gamma_{tt}^k{}_{,k} - \Gamma_{kt}^k{}_{,t} + \Gamma_{tt}^s \Gamma_{ks}^k - \Gamma_{kt}^s \Gamma_{ts}^k \right) \\ &= e^{-2V} \left\{ \partial_t V_t + \partial_r \left(V_r e^{2V} \left(1 - \frac{2M}{r}\right) \right) - \partial_t V_t - \partial_t \left(\frac{M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} \right) \right. \\ &\quad + V_t \left(V_t + \frac{M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} \right) \\ &\quad + V_r e^{2V} \left(1 - \frac{2M}{r}\right) \left(V_r + \left(1 - \frac{2M}{r}\right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) + \frac{2}{r} \right) \\ &\quad \left. - V_t^2 - 2V_r^2 e^{2V} \left(1 - \frac{2M}{r}\right) - \frac{M_t^2}{r^2} \left(1 - \frac{2M}{r}\right)^{-2} \right\} \\ &= e^{-2V} \left\{ V_{rr} e^{2V} \left(1 - \frac{2M}{r}\right) + 2V_r^2 e^{2V} \left(1 - \frac{2M}{r}\right) + V_r e^{2V} \left(\frac{2M}{r^2} - \frac{2M_r}{r} \right) \right. \\ &\quad - \frac{M_{tt}}{r} \left(1 - \frac{2M}{r}\right)^{-1} - \frac{2M_t^2}{r^2} \left(1 - \frac{2M}{r}\right)^{-2} + \frac{V_t M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} \\ &\quad + V_r^2 e^{2V} \left(1 - \frac{2M}{r}\right) + V_r e^{2V} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) + \frac{2V_r e^{2V}}{r} \left(1 - \frac{2M}{r}\right) \\ &\quad \left. - 2V_r^2 e^{2V} \left(1 - \frac{2M}{r}\right) - \frac{M_t^2}{r^2} \left(1 - \frac{2M}{r}\right)^{-2} \right\} \end{aligned}$$

$$\begin{aligned} \text{Ric}(\nu_t, \nu_t) = & \left(V_{rr} + V_r^2 + \frac{2V_r}{r} \right) \left(1 - \frac{2M}{r} \right) + \frac{V_r}{r} \left(\frac{M}{r} - M_r \right) \\ & + \frac{V_t M_t - M_{tt}}{r e^{2V}} \left(1 - \frac{2M}{r} \right)^{-1} - \frac{3M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r} \right)^{-2} \end{aligned} \quad (82)$$

$$\begin{aligned} \text{Ric}(\nu_t, \nu_r) = & e^{-V} \sqrt{1 - \frac{2M}{r}} \left(\Gamma_{tr}^k{}_{,k} - \Gamma_{kr}^k{}_{,t} + \Gamma_{tr}^s \Gamma_{ks}^k - \Gamma_{kr}^s \Gamma_{ts}^k \right) \\ = & e^{-V} \sqrt{1 - \frac{2M}{r}} \left\{ \partial_t(V_r) + \partial_r \left(\frac{M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1} \right) - \partial_t(V_r) \right. \\ & - \partial_t \left(\left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \right) - 2\partial_t \left(\frac{1}{r} \right) + V_r \left(V_t + \frac{M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1} \right) \\ & + \frac{M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1} \left(V_r + \left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) + \frac{2}{r} \right) \\ & \left. - V_r V_t - \frac{2M_t V_r}{r} \left(1 - \frac{2M}{r} \right)^{-1} - \frac{M_t}{r} \left(1 - \frac{2M}{r} \right)^{-2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \right\} \\ = & e^{-V} \sqrt{1 - \frac{2M}{r}} \left\{ \partial_r \partial_t \left(-\frac{1}{2} \log \left(1 - \frac{2M}{r} \right) \right) - \partial_t \partial_r \left(-\frac{1}{2} \log \left(1 - \frac{2M}{r} \right) \right) \right. \\ & \left. + \frac{2M_t}{r^2} \left(1 - \frac{2M}{r} \right)^{-1} \right\} \\ \text{Ric}(\nu_t, \nu_r) = & \frac{2M_t}{r^2 e^V} \left(1 - \frac{2M}{r} \right)^{-1/2} \end{aligned} \quad (83)$$

$$\begin{aligned} \text{Ric}(\nu_r, \nu_r) = & \left(1 - \frac{2M}{r} \right) \left(\Gamma_{rr}^k{}_{,k} - \Gamma_{kr}^k{}_{,r} + \Gamma_{rr}^s \Gamma_{ks}^k - \Gamma_{kr}^s \Gamma_{rs}^k \right) \\ = & \left(1 - \frac{2M}{r} \right) \left\{ \partial_t \left(\frac{M_t}{r e^{2V}} \left(1 - \frac{2M}{r} \right)^{-2} \right) + \partial_r \left(\left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \right) \right. \\ & - \partial_r V_r - \partial_r \left(\left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \right) - 2\partial_r \left(\frac{1}{r} \right) + \frac{V_t M_t}{r e^{2V}} \left(1 - \frac{2M}{r} \right)^{-2} \\ & + \frac{M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r} \right)^{-3} + \left(V_r + \frac{2}{r} \right) \left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \\ & + \left(1 - \frac{2M}{r} \right)^{-2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right)^2 - V_r^2 - \frac{2M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r} \right)^{-3} - \frac{2}{r^2} \\ & \left. - \left(1 - \frac{2M}{r} \right)^{-2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right)^2 \right\} \end{aligned}$$

$$\begin{aligned}
 &= \left(1 - \frac{2M}{r}\right) \left\{ \frac{M_{tt}}{re^{2V}} \left(1 - \frac{2M}{r}\right)^{-2} - \frac{2V_t M_t}{re^{2V}} \left(1 - \frac{2M}{r}\right)^{-2} + \frac{4M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r}\right)^{-3} \right. \\
 &\quad - V_{rr} + \frac{2}{r^2} + \frac{V_t M_t}{re^{2V}} \left(1 - \frac{2M}{r}\right)^{-2} - \frac{M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r}\right)^{-3} - V_r^2 - \frac{2}{r^2} \\
 &\quad \left. + \left(V_r + \frac{2}{r}\right) \left(1 - \frac{2M}{r}\right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2}\right) \right\} \\
 \text{Ric}(\nu_r, \nu_r) &= -(V_{rr} + V_r^2) \left(1 - \frac{2M}{r}\right) - \left(\frac{2}{r^2} + \frac{V_r}{r}\right) \left(\frac{M}{r} - M_r\right) \\
 &\quad - \frac{V_t M_t - M_{tt}}{re^{2V}} \left(1 - \frac{2M}{r}\right)^{-1} + \frac{3M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r}\right)^{-2} \tag{84}
 \end{aligned}$$

$$\begin{aligned}
 \text{Ric}(\nu_\theta, \nu_\theta) &= \frac{1}{r^2} (\Gamma_{\theta\theta}^k{}_{,k} - \Gamma_{k\theta}^k{}_{,\theta} + \Gamma_{\theta\theta}^s \Gamma_{ks}^k - \Gamma_{k\theta}^s \Gamma_{\theta s}^k) \\
 &= \frac{1}{r^2} \left\{ \partial_r \left(-r \left(1 - \frac{2M}{r}\right)\right) - \partial_\theta \left(\frac{\cos \theta}{\sin \theta}\right) - r \left(1 - \frac{2M}{r}\right) \left(V_r + \frac{2}{r}\right) \right. \\
 &\quad \left. - r \left(\frac{M_r}{r} - \frac{M}{r^2}\right) + 2 \left(1 - \frac{2M}{r}\right) - \frac{\cos^2 \theta}{\sin^2 \theta} \right\} \\
 &= \frac{1}{r^2} \left\{ - \left(1 - \frac{2M}{r}\right) - r \left(\frac{2M}{r^2} - \frac{2M_r}{r}\right) + \frac{1}{\sin^2 \theta} - r \left(1 - \frac{2M}{r}\right) \left(V_r + \frac{2}{r}\right) \right. \\
 &\quad \left. + r \left(\frac{M}{r^2} - \frac{M_r}{r}\right) + 2 \left(1 - \frac{2M}{r}\right) - \frac{\cos^2 \theta}{\sin^2 \theta} \right\} \\
 &= \frac{1}{r^2} \left\{ - \frac{\sin^2 \theta}{\sin^2 \theta} + \frac{2M}{r} - \frac{M}{r} + M_r + \frac{1}{\sin^2 \theta} - r V_r \left(1 - \frac{2M}{r}\right) \right. \\
 &\quad \left. - 2 \left(1 - \frac{2M}{r}\right) + 2 \left(1 - \frac{2M}{r}\right) - \frac{\cos^2 \theta}{\sin^2 \theta} \right\} \\
 \text{Ric}(\nu_\theta, \nu_\theta) &= \frac{1}{r^2} \left(\frac{M}{r} + M_r\right) - \frac{V_r}{r} \left(1 - \frac{2M}{r}\right) \tag{85}
 \end{aligned}$$

$$\begin{aligned}
 \text{Ric}(\nu_\varphi, \nu_\varphi) &= \frac{1}{r^2 \sin^2 \theta} (\Gamma_{\varphi\varphi}^k{}_{,k} - \Gamma_{k\varphi}^k{}_{,\varphi} + \Gamma_{\varphi\varphi}^s \Gamma_{ks}^k - \Gamma_{k\varphi}^s \Gamma_{\varphi s}^k) \\
 &= \frac{1}{r^2 \sin^2 \theta} \left\{ \partial_r \left(-r \sin^2 \theta \left(1 - \frac{2M}{r}\right)\right) + \partial_\theta (-\sin \theta \cos \theta) \right. \\
 &\quad - r \sin^2 \theta \left(1 - \frac{2M}{r}\right) \left(V_r + \frac{2}{r}\right) - r \sin^2 \theta \left(\frac{M_r}{r} - \frac{M}{r^2}\right) - \cos^2 \theta \\
 &\quad \left. + 2 \sin^2 \theta \left(1 - \frac{2M}{r}\right) + 2 \cos^2 \theta \right\}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{r^2 \sin^2 \theta} \left\{ -\sin^2 \theta \left(1 - \frac{2M}{r} \right) - r \sin^2 \theta \left(\frac{2M}{r^2} - \frac{2M_r}{r} \right) - \cos^2 \theta + \sin^2 \theta \right. \\
 &\quad \left. - r \sin^2 \theta \left(1 - \frac{2M}{r} \right) \left(V_r + \frac{2}{r} \right) + r \sin^2 \theta \left(\frac{M}{r^2} - \frac{M_r}{r} \right) - \cos^2 \theta \right. \\
 &\quad \left. + 2 \sin^2 \theta \left(1 - \frac{2M}{r} \right) + 2 \cos^2 \theta \right\} \\
 &= \frac{1}{r^2} \left(\frac{2M}{r} - \frac{M}{r} + M_r - r V_r \left(1 - \frac{2M}{r} \right) - 2 \left(1 - \frac{2M}{r} \right) + 2 \left(1 - \frac{2M}{r} \right) \right) \\
 \text{Ric}(\nu_\varphi, \nu_\varphi) &= \frac{1}{r^2} \left(\frac{M}{r} + M_r \right) - \frac{V_r}{r} \left(1 - \frac{2M}{r} \right) = \text{Ric}(\nu_\theta, \nu_\theta). \tag{86}
 \end{aligned}$$

These are the only nonzero components of the Ricci curvature tensor since

$$\begin{aligned}
 \text{Ric}(\nu_t, \nu_\theta) &= \frac{1}{r e^V} \text{Ric}_{t\theta} = \frac{1}{r e^V} \mathbf{R}_{tk\theta}{}^k \\
 &= \frac{1}{r e^V} (\Gamma_{t\theta}{}^k{}_{,k} - \Gamma_{k\theta}{}^k{}_{,t} + \Gamma_{t\theta}{}^s \Gamma_{ks}{}^k - \Gamma_{k\theta}{}^s \Gamma_{ts}{}^k) \\
 &= \frac{1}{r e^V} (-\Gamma_{r\theta}{}^r{}_{,t} - \Gamma_{\theta\theta}{}^\theta{}_{,t} - \Gamma_{\varphi\theta}{}^\varphi{}_{,t} - \Gamma_{k\theta}{}^r \Gamma_{tr}{}^k - \Gamma_{k\theta}{}^\theta \Gamma_{t\theta}{}^k - \Gamma_{k\theta}{}^\varphi \Gamma_{t\varphi}{}^k) \\
 &= \frac{1}{r e^V} (-\Gamma_{\theta\theta}{}^r \Gamma_{tr}{}^\theta - \Gamma_{r\theta}{}^\theta \Gamma_{t\theta}{}^r - \Gamma_{\varphi\theta}{}^\varphi \Gamma_{t\varphi}{}^\varphi) \\
 &= 0 \tag{87}
 \end{aligned}$$

$$\begin{aligned}
 \text{Ric}(\nu_t, \nu_\varphi) &= \frac{1}{r \sin \theta e^V} \text{Ric}_{t\varphi} = \frac{1}{r \sin \theta e^V} \mathbf{R}_{tk\varphi}{}^k \\
 &= \frac{1}{r \sin \theta e^V} (\Gamma_{t\varphi}{}^k{}_{,k} - \Gamma_{k\varphi}{}^k{}_{,t} + \Gamma_{t\varphi}{}^s \Gamma_{ks}{}^k - \Gamma_{k\varphi}{}^s \Gamma_{ts}{}^k) \\
 &= \frac{1}{r \sin \theta e^V} (-\Gamma_{r\varphi}{}^r{}_{,t} - \Gamma_{\theta\varphi}{}^\theta{}_{,t} - \Gamma_{\varphi\varphi}{}^\varphi{}_{,t} - \Gamma_{k\varphi}{}^r \Gamma_{tr}{}^k - \Gamma_{k\varphi}{}^\theta \Gamma_{t\theta}{}^k - \Gamma_{k\varphi}{}^\varphi \Gamma_{t\varphi}{}^k) \\
 &= \frac{1}{r \sin \theta e^V} (-\Gamma_{\varphi\varphi}{}^r \Gamma_{tr}{}^\varphi - \Gamma_{\varphi\varphi}{}^\theta \Gamma_{t\theta}{}^\varphi - \Gamma_{r\varphi}{}^\varphi \Gamma_{t\varphi}{}^r - \Gamma_{\theta\varphi}{}^\varphi \Gamma_{t\varphi}{}^\theta) \\
 &= 0 \tag{88}
 \end{aligned}$$

$$\begin{aligned}
 \text{Ric}(\nu_r, \nu_\theta) &= \frac{1}{r} \sqrt{1 - \frac{2M}{r}} \text{Ric}_{r\theta} = \frac{1}{r} \sqrt{1 - \frac{2M}{r}} \mathbf{R}_{rk\theta}{}^k \\
 &= \frac{1}{r} \sqrt{1 - \frac{2M}{r}} (\Gamma_{r\theta}{}^k{}_{,k} - \Gamma_{k\theta}{}^k{}_{,r} + \Gamma_{r\theta}{}^s \Gamma_{ks}{}^k - \Gamma_{k\theta}{}^s \Gamma_{rs}{}^k) \\
 &= \frac{1}{r} \sqrt{1 - \frac{2M}{r}} (\Gamma_{r\theta}{}^\theta{}_{,\theta} - \Gamma_{k\theta}{}^k{}_{,r} + \Gamma_{r\theta}{}^\theta \Gamma_{k\theta}{}^k - \Gamma_{k\theta}{}^s \Gamma_{rs}{}^k) \\
 &= \frac{1}{r} \sqrt{1 - \frac{2M}{r}} (-\Gamma_{r\theta}{}^r{}_{,r} - \Gamma_{\theta\theta}{}^\theta{}_{,r} - \Gamma_{\varphi\theta}{}^\varphi{}_{,r} + \Gamma_{r\theta}{}^\theta \Gamma_{r\theta}{}^r + \Gamma_{r\theta}{}^\theta \Gamma_{\theta\theta}{}^\theta \\
 &\quad + \Gamma_{r\theta}{}^\theta \Gamma_{\varphi\theta}{}^\varphi - \Gamma_{r\theta}{}^s \Gamma_{rs}{}^r - \Gamma_{\theta\theta}{}^s \Gamma_{rs}{}^\theta - \Gamma_{\varphi\theta}{}^s \Gamma_{rs}{}^\varphi)
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{r} \sqrt{1 - \frac{2M}{r}} \left(\frac{\cos \theta}{r \sin \theta} - \Gamma_{r\theta}{}^\theta \Gamma_{r\theta}{}^r - \Gamma_{\theta\theta}{}^r \Gamma_{rr}{}^\theta - \Gamma_{\varphi\theta}{}^\varphi \Gamma_{r\varphi}{}^\varphi \right) \\
 &= \frac{1}{r} \sqrt{1 - \frac{2M}{r}} \left(\frac{\cos \theta}{r \sin \theta} - \frac{\cos \theta}{r \sin \theta} \right) \\
 &= 0
 \end{aligned} \tag{89}$$

$$\begin{aligned}
 \text{Ric}(\nu_r, \nu_\varphi) &= \frac{1}{r \sin \theta} \sqrt{1 - \frac{2M}{r}} \text{Ric}_{r\varphi} = \frac{1}{r \sin \theta} \sqrt{1 - \frac{2M}{r}} \mathbf{R}_{rk\varphi}{}^k \\
 &= \frac{1}{r \sin \theta} \sqrt{1 - \frac{2M}{r}} (\Gamma_{r\varphi}{}^k{}_{,k} - \Gamma_{k\varphi}{}^k{}_{,r} + \Gamma_{r\varphi}{}^s \Gamma_{ks}{}^k - \Gamma_{k\varphi}{}^s \Gamma_{rs}{}^k) \\
 &= \frac{1}{r \sin \theta} \sqrt{1 - \frac{2M}{r}} (\Gamma_{r\varphi}{}^\varphi{}_{,\varphi} - \Gamma_{k\varphi}{}^k{}_{,r} + \Gamma_{r\varphi}{}^\varphi \Gamma_{k\varphi}{}^k - \Gamma_{k\varphi}{}^s \Gamma_{rs}{}^k) \\
 &= \frac{1}{r \sin \theta} \sqrt{1 - \frac{2M}{r}} (-\Gamma_{r\varphi}{}^r{}_{,r} - \Gamma_{\theta\varphi}{}^\theta{}_{,r} - \Gamma_{\varphi\varphi}{}^\varphi{}_{,r} + \Gamma_{r\varphi}{}^\varphi \Gamma_{r\varphi}{}^r + \Gamma_{r\varphi}{}^\varphi \Gamma_{\theta\varphi}{}^\theta \\
 &\quad + \Gamma_{r\varphi}{}^\varphi \Gamma_{\varphi\varphi}{}^\varphi - \Gamma_{r\varphi}{}^s \Gamma_{rs}{}^r - \Gamma_{\theta\varphi}{}^s \Gamma_{rs}{}^\theta - \Gamma_{\varphi\varphi}{}^s \Gamma_{rs}{}^\varphi) \\
 &= \frac{1}{r \sin \theta} \sqrt{1 - \frac{2M}{r}} (-\Gamma_{r\varphi}{}^\varphi \Gamma_{r\varphi}{}^r - \Gamma_{\theta\varphi}{}^\varphi \Gamma_{r\varphi}{}^\theta - \Gamma_{\varphi\varphi}{}^r \Gamma_{rr}{}^\varphi - \Gamma_{\varphi\varphi}{}^\theta \Gamma_{r\theta}{}^\varphi) \\
 &= 0
 \end{aligned} \tag{90}$$

$$\begin{aligned}
 \text{Ric}(\nu_\theta, \nu_\varphi) &= \frac{1}{r^2 \sin \theta} \text{Ric}_{\theta\varphi} = \frac{1}{r^2 \sin \theta} \mathbf{R}_{\theta k\varphi}{}^k \\
 &= \frac{1}{r^2 \sin \theta} (\Gamma_{\theta\varphi}{}^k{}_{,k} - \Gamma_{k\varphi}{}^k{}_{,\theta} + \Gamma_{\theta\varphi}{}^s \Gamma_{ks}{}^k - \Gamma_{k\varphi}{}^s \Gamma_{\theta s}{}^k) \\
 &= \frac{1}{r^2 \sin \theta} (\Gamma_{\theta\varphi}{}^\varphi{}_{,\varphi} - \Gamma_{k\varphi}{}^k{}_{,\theta} + \Gamma_{\theta\varphi}{}^\varphi \Gamma_{k\varphi}{}^k - \Gamma_{k\varphi}{}^s \Gamma_{\theta s}{}^k) \\
 &= \frac{1}{r^2 \sin \theta} (-\Gamma_{r\varphi}{}^r{}_{,\theta} - \Gamma_{\theta\varphi}{}^\theta{}_{,\theta} - \Gamma_{\varphi\varphi}{}^\varphi{}_{,\theta} + \Gamma_{\theta\varphi}{}^\varphi \Gamma_{r\varphi}{}^r + \Gamma_{\theta\varphi}{}^\varphi \Gamma_{\theta\varphi}{}^\theta \\
 &\quad + \Gamma_{\theta\varphi}{}^\varphi \Gamma_{\varphi\varphi}{}^\varphi - \Gamma_{k\varphi}{}^r \Gamma_{\theta r}{}^k - \Gamma_{k\varphi}{}^\theta \Gamma_{\theta\theta}{}^k - \Gamma_{k\varphi}{}^\varphi \Gamma_{\theta\varphi}{}^k) \\
 &= \frac{1}{r^2 \sin \theta} (-\Gamma_{\theta\varphi}{}^r \Gamma_{\theta r}{}^\theta - \Gamma_{r\varphi}{}^\theta \Gamma_{\theta\theta}{}^r - \Gamma_{\varphi\varphi}{}^\varphi \Gamma_{\theta\varphi}{}^\varphi) \\
 &= 0
 \end{aligned} \tag{91}$$

The other components are 0 by the symmetry of the Ricci curvature tensor. To get the last statement, we note that by the last line above and Lemma A.1-2, we have that

$$R = \sum_{\eta} g(\nu_\eta, \nu_\eta) \text{Ric}(\nu_\eta, \nu_\eta) = -\text{Ric}(\nu_t, \nu_t) + \text{Ric}(\nu_r, \nu_r) + 2 \text{Ric}(\nu_\theta, \nu_\theta). \tag{92}$$

From here, it is a matter of algebra and the statements above to get the desired equation for R . $\square$

Next, we prove Proposition 3.7.

Proof of Proposition 3.7. We already have the equations resulting from two of the components of the Einstein equation, namely (30) and (36), which we reprint here for convenience.

$$M_r = 4\pi r^2 \mu \quad (93)$$

$$V_r = \left(1 - \frac{2M}{r}\right)^{-1} \left(\frac{M}{r^2} + 4\pi r P\right) \quad (94)$$

In order to show that solving these two equations is sufficient to solve the entire Einstein equation, we must show that if these equations hold, so do the other components of the Einstein equation. To do so, we first need to write down the other components of the Einstein equation. Recall that the above equations come from the ν_t, ν_t and ν_r, ν_r components of the Einstein equation. Thus we only have the ν_t, ν_r and ν_θ, ν_θ components left to compute (the ν_φ, ν_φ component is identical to ν_θ, ν_θ component). Then, by Corollary 3.3 and (42), we have the following.

$$\begin{aligned} G(\nu_t, \nu_r) &= 8\pi T(\nu_t, \nu_r) \\ \frac{2M_t}{r^2 e^V} \left(1 - \frac{2M}{r}\right)^{-1/2} &= 8\pi \rho \\ M_t &= 4\pi r^2 e^V \rho \sqrt{1 - \frac{2M}{r}} \end{aligned} \quad (95)$$

and

$$\begin{aligned} G(\nu_\theta, \nu_\theta) &= 8\pi T(\nu_\theta, \nu_\theta) \\ 8\pi Q &= \left(V_{rr} + V_r^2 + \frac{V_r}{r}\right) \left(1 - \frac{2M}{r}\right) + \left(\frac{M}{r} - M_r\right) \left(\frac{1}{r^2} + \frac{V_r}{r}\right) \\ &\quad + \frac{V_t M_t - M_{tt}}{r e^{2V}} \left(1 - \frac{2M}{r}\right)^{-1} - \frac{3M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r}\right)^{-2} \end{aligned} \quad (96)$$

Next, since T is a stress-energy tensor, we know that it is divergence free, that is, $\nabla_g \cdot T = 0$, where $\nabla_g \cdot T$ denotes the divergence of the tensor T with respect to the metric g . We will compute next what this equation means in terms of the functions μ , P , ρ , and Q . In what follows, the summing indices run through the set $\{t, r, \theta, \varphi\}$ and we will use the Einstein summation convention wherever it applies, while specifically denoting any other summations of a different form. First, we define

$$\varepsilon_k = g(\nu_k, \nu_k) \quad (97)$$

for all $k \in \{t, r, \theta, \varphi\}$. Then we can write the divergence of T as follows.

$$\begin{aligned} \nabla_g \cdot T &= \sum_k \varepsilon_k (\nabla_{\nu_k} T)(\nu_k, \partial_j) dx^j \\ &= -(\nabla_{\nu_t} T)(\nu_t, \partial_j) dx^j + (\nabla_{\nu_r} T)(\nu_r, \partial_j) dx^j + (\nabla_{\nu_\theta} T)(\nu_\theta, \partial_j) dx^j + (\nabla_{\nu_\varphi} T)(\nu_\varphi, \partial_j) dx^j \end{aligned} \quad (98)$$

We will simplify each of these four terms individually. By equations (81), (22), and (42), we have that

$$\begin{aligned} (\nabla_{\nu_t} T)(\nu_t, \partial_k) dx^k &= e^{-2V} (\nabla_t T)(\partial_t, \partial_k) dx^k \\ &= e^{-2V} \left[\partial_t (T(\partial_t, \partial_k)) - \Gamma_{tt}^m T(\partial_m, \partial_k) - \Gamma_{tk}^m T(\partial_t, \partial_m) \right] dx^k \end{aligned}$$

$$\begin{aligned}
 &= e^{-2V} \left[\partial_t(T(\partial_t, \partial_t)) dt + \partial_t(T(\partial_t, \partial_r)) dr - 2\Gamma_{tt}^t T(\partial_t, \partial_t) dt - 2\Gamma_{tt}^r T(\partial_t, \partial_r) dt \right. \\
 &\quad \left. - (\Gamma_{tt}^t + \Gamma_{tr}^r) T(\partial_t, \partial_r) dr - \Gamma_{tr}^t T(\partial_t, \partial_t) dr - \Gamma_{tt}^r T(\partial_r, \partial_r) dr \right] \\
 &= e^{-2V} \left[\partial_t(e^{2V} \mu) dt + \partial_t \left(e^V \rho \left(1 - \frac{2M}{r} \right)^{-1/2} \right) dr - 2V_t e^{2V} \mu dt \right. \\
 &\quad \left. - 2\rho V_r e^{3V} \sqrt{1 - \frac{2M}{r}} dt - \left(V_t + \frac{M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1} \right) e^V \rho \left(1 - \frac{2M}{r} \right)^{-1/2} dr \right. \\
 &\quad \left. - V_r e^{2V} \mu dr - V_r e^{2V} P dr \right] \\
 &= e^{-2V} \left[\left(2V_t e^{2V} \mu + e^{2V} \mu_t - 2V_t e^{2V} \mu - 2\rho V_r e^{3V} \sqrt{1 - \frac{2M}{r}} \right) dt \right. \\
 &\quad \left. + \left(V_t e^V \rho \left(1 - \frac{2M}{r} \right)^{-1/2} + e^V \rho_t \left(1 - \frac{2M}{r} \right)^{-1/2} + \frac{e^V \rho M_t}{r} \left(1 - \frac{2M}{r} \right)^{-3/2} \right. \right. \\
 &\quad \left. \left. - \left(V_t + \frac{M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1} \right) e^V \rho \left(1 - \frac{2M}{r} \right)^{-1/2} - V_r e^{2V} \mu - V_r e^{2V} P \right) dr \right] \\
 (\nabla_{\nu_t} T)(\nu_t, \partial_k) dx^k &= \left(\mu_t - 2\rho V_r e^V \sqrt{1 - \frac{2M}{r}} \right) dt + \left(\rho_t e^{-V} \left(1 - \frac{2M}{r} \right)^{-1/2} - V_r (\mu + P) \right) dr \\
 &\hspace{15em} (99)
 \end{aligned}$$

Next, we have

$$\begin{aligned}
 (\nabla_{\nu_r} T)(\nu_r, \partial_k) dx^k &= \left(1 - \frac{2M}{r} \right) (\nabla_r T)(\partial_r, \partial_k) dx^k \\
 &= \left(1 - \frac{2M}{r} \right) \left[\partial_r(T(\partial_r, \partial_k)) - \Gamma_{rr}^m T(\partial_m, \partial_k) - \Gamma_{rk}^m T(\partial_r, \partial_m) \right] dx^k \\
 &= \left(1 - \frac{2M}{r} \right) \left[\partial_r(T(\partial_r, \partial_t)) dt + \partial_r(T(\partial_r, \partial_r)) dr - (\Gamma_{rr}^r + \Gamma_{rt}^t) T(\partial_t, \partial_r) dt \right. \\
 &\quad \left. - 2\Gamma_{rr}^t T(\partial_t, \partial_r) dr - 2\Gamma_{rr}^r T(\partial_r, \partial_r) dr - \Gamma_{rr}^t T(\partial_t, \partial_t) dt - \Gamma_{rt}^r T(\partial_r, \partial_r) dt \right] \\
 &= \left(1 - \frac{2M}{r} \right) \left[\partial_r \left(\rho e^V \left(1 - \frac{2M}{r} \right)^{-1/2} \right) dt + \partial_r \left(P \left(1 - \frac{2M}{r} \right)^{-1} \right) dr \right. \\
 &\quad \left. - \rho e^V \left(1 - \frac{2M}{r} \right)^{-1/2} \left[\left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) + V_r \right] dt \right. \\
 &\quad \left. - \frac{2\rho M_t}{r e^V} \left(1 - \frac{2M}{r} \right)^{-5/2} dr - 2P \left(1 - \frac{2M}{r} \right)^{-2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) dr \right. \\
 &\quad \left. - \frac{\mu M_t}{r} \left(1 - \frac{2M}{r} \right)^{-2} dt - \frac{P M_t}{r} \left(1 - \frac{2M}{r} \right)^{-2} dt \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \left(1 - \frac{2M}{r}\right) \left[\left(\rho e^V \left(1 - \frac{2M}{r}\right)^{-3/2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) + \rho V_r e^V \left(1 - \frac{2M}{r}\right)^{-1/2} \right. \right. \\
 &\quad \left. \left. + \rho_r e^V \left(1 - \frac{2M}{r}\right)^{-1/2} - \frac{\mu M_t}{r} \left(1 - \frac{2M}{r}\right)^{-2} - \frac{P M_t}{r} \left(1 - \frac{2M}{r}\right)^{-2} \right. \right. \\
 &\quad \left. \left. - \rho e^V \left(1 - \frac{2M}{r}\right)^{-1/2} \left[\left(1 - \frac{2M}{r}\right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) + V_r \right] \right) dt \right. \\
 &\quad \left. + \left(P_r \left(1 - \frac{2M}{r}\right)^{-1} + 2P \left(1 - \frac{2M}{r}\right)^{-2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \right. \right. \\
 &\quad \left. \left. - \frac{2\rho M_t}{r e^V} \left(1 - \frac{2M}{r}\right)^{-5/2} - 2P \left(1 - \frac{2M}{r}\right)^{-2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \right) dr \right] \\
 (\nabla_{\nu_r} T)(\nu_r, \partial_k) dx^k &= \left(\rho_r e^V \sqrt{1 - \frac{2M}{r}} - \frac{M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} (\mu + P) \right) dt \\
 &\quad + \left(P_r - \frac{2\rho M_t}{r e^V} \left(1 - \frac{2M}{r}\right)^{-3/2} \right) dr
 \end{aligned} \tag{100}$$

Thirdly,

$$\begin{aligned}
 (\nabla_{\nu_\theta} T)(\nu_\theta, \partial_k) dx^k &= \frac{1}{r^2} (\nabla_\theta T)(\partial_\theta, \partial_k) dx^k \\
 &= \frac{1}{r^2} \left[\partial_\theta (T(\partial_\theta, \partial_k)) - \Gamma_{\theta\theta}^m T(\partial_m, \partial_k) - \Gamma_{\theta k}^m T(\partial_\theta, \partial_m) \right] dx^k \\
 &= \frac{1}{r^2} \left[-\Gamma_{\theta\theta}^r T(\partial_r, \partial_t) dt - \Gamma_{\theta\theta}^r T(\partial_r, \partial_r) dr - \Gamma_{\theta r}^\theta T(\partial_\theta, \partial_\theta) dr \right] \\
 &= \frac{1}{r^2} \left[r \rho e^V \sqrt{1 - \frac{2M}{r}} dt + r P dr - r Q dr \right] \\
 (\nabla_{\nu_\theta} T)(\nu_\theta, \partial_k) dx^k &= \frac{\rho e^V}{r} \sqrt{1 - \frac{2M}{r}} dt + \frac{P - Q}{r} dr
 \end{aligned} \tag{101}$$

Lastly, we have that

$$\begin{aligned}
 (\nabla_{\nu_\varphi} T)(\nu_\varphi, \partial_k) dx^k &= \frac{1}{r^2 \sin^2 \theta} (\nabla_\varphi T)(\partial_\varphi, \partial_k) dx^k \\
 &= \frac{1}{r^2 \sin^2 \theta} \left[\partial_\varphi (T(\partial_\varphi, \partial_k)) - \Gamma_{\varphi\varphi}^m T(\partial_m, \partial_k) - \Gamma_{\varphi k}^m T(\partial_\varphi, \partial_m) \right] dx^k \\
 &= \frac{1}{r^2 \sin^2 \theta} \left[-\Gamma_{\varphi\varphi}^r T(\partial_r, \partial_t) dt - \Gamma_{\varphi\varphi}^r T(\partial_r, \partial_r) dr - \Gamma_{\varphi\varphi}^\theta T(\partial_\theta, \partial_\theta) d\theta \right. \\
 &\quad \left. - \Gamma_{\varphi r}^\varphi T(\partial_\varphi, \partial_\varphi) dr - \Gamma_{\varphi\theta}^\varphi T(\partial_\varphi, \partial_\varphi) d\theta \right] \\
 &= \frac{1}{r^2 \sin^2 \theta} \left[r \rho e^V \sin^2 \theta \sqrt{1 - \frac{2M}{r}} dt + r \sin^2 \theta P dr \right. \\
 &\quad \left. + r^2 Q \sin \theta \cos \theta d\theta - r Q \sin^2 \theta dr - r^2 Q \cos \theta \sin \theta d\theta \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{\rho e^V}{r} \sqrt{1 - \frac{2M}{r}} dt + \frac{P - Q}{r} dr \\
 (\nabla_{\nu_\varphi} T)(\nu_\varphi, \partial_k) dx^k &= (\nabla_{\nu_\theta} T)(\nu_\theta, \partial_k) dx^k
 \end{aligned} \tag{102}$$

Then equation (98) becomes

$$\begin{aligned}
 \nabla_g \cdot T &= -(\nabla_{\nu_t} T)(\nu_t, \partial_j) dx^j + (\nabla_{\nu_r} T)(\nu_r, \partial_j) dx^j + (\nabla_{\nu_\theta} T)(\nu_\theta, \partial_j) dx^j + (\nabla_{\nu_\varphi} T)(\nu_\varphi, \partial_j) dx^j \\
 &= -\left(\mu_t - 2\rho V_r e^V \sqrt{1 - \frac{2M}{r}}\right) dt - \left(\rho_t e^{-V} \left(1 - \frac{2M}{r}\right)^{-1/2} - V_r(\mu + P)\right) dr \\
 &\quad + \left(\rho_r e^V \sqrt{1 - \frac{2M}{r}} - \frac{M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} (\mu + P)\right) dt \\
 &\quad + \left(P_r - \frac{2\rho M_t}{r e^V} \left(1 - \frac{2M}{r}\right)^{-3/2}\right) dr + \frac{2\rho e^V}{r} \sqrt{1 - \frac{2M}{r}} dt + \frac{2(P - Q)}{r} dr \\
 &= \left(-\mu_t + 2\rho e^V \sqrt{1 - \frac{2M}{r}} \left(V_r + \frac{1}{r}\right) + \rho_r e^V \sqrt{1 - \frac{2M}{r}} - \frac{M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} (\mu + P)\right) dt \\
 &\quad + \left(-\rho_t e^{-V} \left(1 - \frac{2M}{r}\right)^{-1/2} + V_r(\mu + P) + P_r - \frac{2\rho M_t}{r e^V} \left(1 - \frac{2M}{r}\right)^{-3/2} + \frac{2(P - Q)}{r}\right) dr
 \end{aligned} \tag{103}$$

Then $\nabla_g \cdot T = 0$ yields the following two equations

$$\mu_t = 2\rho e^V \sqrt{1 - \frac{2M}{r}} \left(V_r + \frac{1}{r}\right) + \rho_r e^V \sqrt{1 - \frac{2M}{r}} - \frac{M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} (\mu + P) \tag{104a}$$

$$\rho_t = e^V \sqrt{1 - \frac{2M}{r}} \left(V_r(\mu + P) + P_r + \frac{2(P - Q)}{r}\right) - \frac{2\rho M_t}{r} \left(1 - \frac{2M}{r}\right)^{-1} \tag{104b}$$

We are now ready to show that solving (30) and (36) on each $t = \text{constant}$ slice also solves the Einstein equation. In other words, as stated earlier, we want to show that solving these two equations for each constant t automatically satisfies the remaining two unique nonzero components of the Einstein equation, (95) and (96). We will show first that (30), (36), and (104) implies that (95) holds as well. To do this, we first need to show that (30) and (95) are compatible, that is, that, given these equations, $M_{rt} = M_{tr}$. Differentiating (30) with respect to t yields,

$$\begin{aligned}
 \partial_t M_r &= \partial_t (4\pi r^2 \mu) \\
 M_{rt} &= 4\pi r^2 \mu_t
 \end{aligned} \tag{105}$$

while differentiating (95) with respect to r yields,

$$\begin{aligned}\partial_r M_t &= \partial_r \left(4\pi r^2 e^V \rho \sqrt{1 - \frac{2M}{r}} \right) \\ M_{tr} &= 4\pi \left[2re^V \rho \sqrt{1 - \frac{2M}{r}} + r^2 V_r e^V \rho \sqrt{1 - \frac{2M}{r}} \right. \\ &\quad \left. + r^2 e^V \rho_r \sqrt{1 - \frac{2M}{r}} - r^2 e^V \rho \left(1 - \frac{2M}{r} \right)^{-1/2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \right]\end{aligned}$$

To make the simplification process easier, to the right hand side above, we will add in and subtract out a copy of the second term in the sum above, namely, the term $4\pi r^2 V_r e^V \rho \sqrt{1 - 2M/r}$.

$$\begin{aligned}M_{tr} &= 4\pi \left[2re^V \rho \sqrt{1 - \frac{2M}{r}} + 2r^2 V_r e^V \rho \sqrt{1 - \frac{2M}{r}} - r^2 V_r e^V \rho \sqrt{1 - \frac{2M}{r}} \right. \\ &\quad \left. + r^2 e^V \rho_r \sqrt{1 - \frac{2M}{r}} - r^2 e^V \rho \left(1 - \frac{2M}{r} \right)^{-1/2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \right] \\ &= 4\pi r^2 \left[2\rho e^V \sqrt{1 - \frac{2M}{r}} \left(V_r + \frac{1}{r} \right) + \rho_r e^V \sqrt{1 - \frac{2M}{r}} \right. \\ &\quad \left. - V_r e^V \rho \sqrt{1 - \frac{2M}{r}} - e^V \rho \left(1 - \frac{2M}{r} \right)^{-1/2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \right] \\ &= 4\pi r^2 \left[\mu_t + \frac{M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1} (\mu + P) \right. \\ &\quad \left. - e^V \rho \left(1 - \frac{2M}{r} \right)^{-1/2} \left(\frac{M}{r^2} + 4\pi r P \right) - e^V \rho \left(1 - \frac{2M}{r} \right)^{-1/2} \left(4\pi r \mu - \frac{M}{r^2} \right) \right] \\ &= 4\pi r^2 \left[\mu_t + \frac{M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1} (\mu + P) - 4\pi r \rho e^V \left(1 - \frac{2M}{r} \right)^{-1/2} (\mu + P) \right] \\ &= 4\pi r^2 \mu_t\end{aligned}\tag{106}$$

where in the next to last line we made substitutions using equations (30), (36), and (104a) and the last line used (95). Thus if $\nabla_g \cdot T = 0$, (30) and (95) are compatible, that is, $M_{tr} = M_{rt}$ everywhere the equations are defined, that is, wherever $r \neq 0$. However, by using L'Hôpital's rule on the equations above coupled with the fact that at $r = 0$, $M = M_r = M_{rr} = 0$ for all t , the equation $M_{tr} = M_{rt}$ still holds at the central value $r = 0$. This implies that there exists a function $M(t, r)$ which satisfies both (30) and (36) everywhere, which of course is the metric function we seek.

If, for all values of t , we integrate (30) with the initial condition $M = 0$ at $r = 0$, we will obtain some function $M^*(t, r)$ which satisfies (30) everywhere. Then we have that $M_r^* = M_r$ everywhere, where M is the function that satisfies both (30) and (36). This implies that $M^*(t, r) = M(t, r) + f(t)$ for some smooth function $f(t)$. However, we also have that $M^*(t, 0) =$

$M(t, 0) = 0$ for all t , which implies that $f(t) \equiv 0$. Hence $M^*(t, r) = M(t, r)$. Thus the function we obtain by integrating (30) with compatible initial conditions necessarily also satisfies (95).

Finally, we use equations (30), (36), (104), and (95) (since the first three imply it) to show that (96) is automatically satisfied. In order to do this, we will have to compute V_{rr} and M_{tt} . They are

$$\begin{aligned} \partial_t M_t &= \partial_t \left(4\pi r^2 e^V \rho \sqrt{1 - \frac{2M}{r}} \right) \\ M_{tt} &= 4\pi r^2 \left[V_t e^V \rho \sqrt{1 - \frac{2M}{r}} + e^V \rho_t \sqrt{1 - \frac{2M}{r}} - \frac{e^V \rho M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1/2} \right] \\ &= V_t M_t + 4\pi r^2 e^V \left[\rho_t \sqrt{1 - \frac{2M}{r}} - \frac{\rho M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1/2} \right] \end{aligned} \quad (107)$$

and

$$\begin{aligned} \partial_r V_r &= \partial_r \left[\left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M}{r^2} + 4\pi r P \right) \right] \\ V_{rr} &= 2 \left(1 - \frac{2M}{r} \right)^{-2} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) \left(\frac{M}{r^2} + 4\pi r P \right) + \left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r^2} - \frac{2M}{r^3} + 4\pi P + 4\pi r P_r \right) \\ &= 2V_r \left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) + \left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r^2} - \frac{M}{r^3} - \frac{M}{r^3} + 4\pi P + 4\pi r P_r \right) \\ &= 2V_r \left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r} - \frac{M}{r^2} \right) - \frac{V_r}{r} + \left(1 - \frac{2M}{r} \right)^{-1} \left(\frac{M_r}{r^2} - \frac{M}{r^3} + 8\pi P + 4\pi r P_r \right) \\ V_{rr} &= \left(2\frac{V_r}{r} + \frac{1}{r^2} \right) \left(1 - \frac{2M}{r} \right)^{-1} \left(M_r - \frac{M}{r} \right) - \frac{V_r}{r} + 4\pi \left(1 - \frac{2M}{r} \right)^{-1} (2P + rP_r) \end{aligned} \quad (108)$$

Making these substitutions into (96) yields

$$\begin{aligned} 8\pi Q &= \left(V_{rr} + V_r^2 + \frac{V_r}{r} \right) \left(1 - \frac{2M}{r} \right) + \left(\frac{M}{r} - M_r \right) \left(\frac{1}{r^2} + \frac{V_r}{r} \right) \\ &\quad + \frac{V_t M_t - M_{tt}}{r e^{2V}} \left(1 - \frac{2M}{r} \right)^{-1} - \frac{3M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r} \right)^{-2} \\ &= 4\pi (2P + rP_r) + V_r^2 \left(1 - \frac{2M}{r} \right) + \frac{V_r}{r} \left(M_r - \frac{M}{r} \right) \\ &\quad - \frac{4\pi r}{e^V} \left[\rho_t \left(1 - \frac{2M}{r} \right)^{-1/2} - \frac{\rho M_t}{r} \left(1 - \frac{2M}{r} \right)^{-3/2} \right] - \frac{3M_t^2}{r^2 e^{2V}} \left(1 - \frac{2M}{r} \right)^{-2} \\ &= 4\pi (2P + rP_r) + V_r \left(\frac{M}{r^2} + 4\pi r P \right) + \frac{V_r}{r} \left(M_r - \frac{M}{r} \right) \\ &\quad - \frac{4\pi r}{e^V} \left[\rho_t \left(1 - \frac{2M}{r} \right)^{-1/2} - \frac{\rho M_t}{r} \left(1 - \frac{2M}{r} \right)^{-3/2} \right] - \frac{12\pi \rho M_t}{e^V} \left(1 - \frac{2M}{r} \right)^{-3/2} \end{aligned}$$

$$\begin{aligned}
 &= 4\pi(2P + rP_r) + 4\pi rV_rP + \frac{V_rM_r}{r} - \frac{4\pi r\rho_t}{e^V} \left(1 - \frac{2M}{r}\right)^{-1/2} - \frac{8\pi\rho M_t}{e^V} \left(1 - \frac{2M}{r}\right)^{-3/2} \\
 &= 4\pi(2P + rP_r) + 4\pi rV_r(P + \mu) - \frac{4\pi r\rho_t}{e^V} \left(1 - \frac{2M}{r}\right)^{-1/2} - \frac{8\pi\rho M_t}{e^V} \left(1 - \frac{2M}{r}\right)^{-3/2} \\
 &= 4\pi(2P + rP_r) + 4\pi rV_r(P + \mu) - \frac{8\pi\rho M_t}{e^V} \left(1 - \frac{2M}{r}\right)^{-3/2} \\
 &\quad - 4\pi r \left(V_r(\mu + P) + P_r + \frac{2(P - Q)}{r} \right) + \frac{8\pi\rho M_t}{e^V} \left(1 - \frac{2M}{r}\right)^{-3/2} \\
 &= 8\pi Q
 \end{aligned} \tag{109}$$

where the last two lines follow from using (104b). Thus so long as equation (104) holds, equations (30), (36), and (95) imply equation (96). Since equations (104), (30), and (36) imply equation (95) and equation (104) follows from $\nabla_g \cdot T = 0$, we have that $\nabla_g \cdot T = 0$ implies that solving equations (30) and (36) at each time t is sufficient to solve the entire Einstein equation, which was the desired result. $\square$

The proof of Lemma 4.1 follows.

Proof of Lemma 4.1. Recall equation (47). Then to compute the components of T , we first have that

$$df = f_t dt + f_r dr = pe^V \sqrt{1 - \frac{2M}{r}} dt + f_r dr \tag{110}$$

$$d\bar{f} = \bar{f}_t dt + \bar{f}_r dr = \bar{p}e^V \sqrt{1 - \frac{2M}{r}} dt + \bar{f}_r dr. \tag{111}$$

Then, since g is diagonal, this implies that

$$\begin{aligned}
 |df|^2 &= g(df, d\bar{f}) = g \left(pe^V \sqrt{1 - \frac{2M}{r}} dt + f_r dr, \bar{p}e^V \sqrt{1 - \frac{2M}{r}} dt + \bar{f}_r dr \right) \\
 &= e^{2V} \left(1 - \frac{2M}{r} \right) p\bar{p} g(dt, dt) + f_r \bar{f}_r g(dr, dr) \\
 &= \left(1 - \frac{2M}{r} \right) (|f_r|^2 - |p|^2).
 \end{aligned} \tag{112}$$

With these facts, we now compute the quantities in question and do so in the same order as they were presented. Thus we have the following.

$$\begin{aligned}
 T(\nu_t, \nu_t) &= \mu_0 \left(\frac{2}{\Upsilon^2} df(\nu_t) d\bar{f}(\nu_t) - \left[\left(1 - \frac{2M}{r} \right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} + |f|^2 \right] g(\nu_t, \nu_t) \right) \\
 &= \mu_0 \left(\frac{2}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) p\bar{p} + |f|^2 + \left(1 - \frac{2M}{r} \right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} \right) \\
 &= \mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r} \right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right).
 \end{aligned} \tag{113}$$

Next, because ν_η is an everywhere orthonormal basis,

$$\begin{aligned} T(\nu_t, \nu_r) &= \mu_0 \left(\frac{1}{\Upsilon^2} [df(\nu_t)d\bar{f}(\nu_r) + d\bar{f}(\nu_t)df(\nu_r)] - \left[\left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} + |f|^2 \right] g(\nu_t, \nu_r) \right) \\ &= \mu_0 \left(\frac{1}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) (p\bar{f}_r + f_r\bar{p}) \right) \\ &= \frac{2\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \text{Re}(f_r\bar{p}). \end{aligned} \quad (114)$$

$$\begin{aligned} T(\nu_r, \nu_r) &= \mu_0 \left(\frac{2}{\Upsilon^2} df(\nu_r)d\bar{f}(\nu_r) - \left[\left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} + |f|^2 \right] g(\nu_r, \nu_r) \right) \\ &= \mu_0 \left(\frac{2}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) f_r\bar{f}_r - \left[\left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} + |f|^2 \right] \right) \\ &= \mu_0 \left(-|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right). \\ T(\nu_\theta, \nu_\theta) &= \mu_0 \left(\frac{2}{\Upsilon^2} df(\nu_\theta)d\bar{f}(\nu_\theta) - \left[\left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} + |f|^2 \right] g(\nu_\theta, \nu_\theta) \right) \\ &= -\mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} \right). \end{aligned} \quad (115)$$

Finally,

$$\begin{aligned} T(\nu_\varphi, \nu_\varphi) &= \mu_0 \left(\frac{2}{\Upsilon^2} df(\nu_\varphi)d\bar{f}(\nu_\varphi) - \left[\left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} + |f|^2 \right] g(\nu_\varphi, \nu_\varphi) \right) \\ &= -\mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} \right) \end{aligned} \quad (116)$$

so that $T(\nu_\varphi, \nu_\varphi) = T(\nu_\theta, \nu_\theta)$. □

Finally, we prove Lemma 4.4.

Proof of Lemma 4.4. By equation (42) and Lemma 4.1, we have that

$$\mu = \mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \quad (117a)$$

$$\rho = \frac{2\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \text{Re}(f_r\bar{p}) \quad (117b)$$

$$P = \mu_0 \left(-|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \quad (117c)$$

$$Q = -\mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r}\right) \frac{|f_r|^2 - |p|^2}{\Upsilon^2} \right) \quad (117d)$$

We know from a previous proof that $\nabla_g \cdot T = 0$ is equivalent to equation (104). Moreover, by (48) and Lemma 4.3, we have that (46) is equivalent to (54) and (58). Thus it suffices to show that given (117), equations (54) and (58) imply equation (104).

We will start with (104a). On one hand, the left hand side is equal to differentiating (117a) with respect to t .

$$\begin{aligned}
 \mu_t &= \partial_t \left[\mu_0 \left(|f|^2 + \left(1 - \frac{2M}{r} \right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \right] \\
 &= \mu_0 \left[f_t \bar{f} + f \bar{f}_t - \frac{2M_t}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) + \left(1 - \frac{2M}{r} \right) \frac{f_{rt} \bar{f}_r + f_r \bar{f}_{rt} + p_t \bar{p} + p \bar{p}_t}{\Upsilon^2} \right] \\
 &= \mu_0 \left[e^V \sqrt{1 - \frac{2M}{r}} (p \bar{f} + f \bar{p}) - \frac{2M_t}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) + \left(1 - \frac{2M}{r} \right) \frac{f_{rt} \bar{f}_r + f_r \bar{f}_{rt} + p_t \bar{p} + p \bar{p}_t}{\Upsilon^2} \right] \\
 \mu_t &= \mu_0 \left[\bar{p} \left(f e^V \sqrt{1 - \frac{2M}{r}} + \frac{p_t}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \right) + p \left(\bar{f} e^V \sqrt{1 - \frac{2M}{r}} + \frac{\bar{p}_t}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \right) \right. \\
 &\quad \left. - \frac{2M_t}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) + \left(1 - \frac{2M}{r} \right) \frac{\bar{f}_r \partial_r (p e^V \sqrt{1 - \frac{2M}{r}}) + f_r \partial_r (\bar{p} e^V \sqrt{1 - \frac{2M}{r}})}{\Upsilon^2} \right] \tag{118a}
 \end{aligned}$$

On the other hand, we can substitute (117) into (104a).

$$\begin{aligned}
 \mu_t &= 2\rho e^V \sqrt{1 - \frac{2M}{r}} \left(V_r + \frac{1}{r} \right) + \rho_r e^V \sqrt{1 - \frac{2M}{r}} - \frac{M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1} (\mu + P) \\
 &= 2e^V \sqrt{1 - \frac{2M}{r}} \left(\frac{2\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \operatorname{Re}(f_r \bar{p}) \right) \left(V_r + \frac{1}{r} \right) \\
 &\quad + \partial_r \left[\frac{2\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \operatorname{Re}(f_r \bar{p}) \right] e^V \sqrt{1 - \frac{2M}{r}} - \frac{2M_t \mu_0}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \\
 &= \frac{2\mu_0}{\Upsilon^2} e^V \left(1 - \frac{2M}{r} \right)^{3/2} (f_r \bar{p} + p \bar{f}_r) \left(V_r + \frac{1}{r} \right) \\
 &\quad + \frac{\mu_0}{\Upsilon^2} \partial_r \left[\left(1 - \frac{2M}{r} \right) (f_r \bar{p} + p \bar{f}_r) \right] e^V \sqrt{1 - \frac{2M}{r}} - \frac{2M_t \mu_0}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \\
 &= \bar{p} \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \left[f_r V_r e^V \sqrt{1 - \frac{2M}{r}} + \frac{2f_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] \\
 &\quad + p \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \left[\bar{f}_r V_r e^V \sqrt{1 - \frac{2M}{r}} + \frac{2\bar{f}_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] \\
 &\quad + \frac{\mu_0 V_r e^V}{\Upsilon^2} \left(1 - \frac{2M}{r} \right)^{3/2} (f_r \bar{p} + p \bar{f}_r) - \frac{2M_t \mu_0}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \\
 &\quad + \frac{\mu_0 e^V}{\Upsilon^2} \sqrt{1 - \frac{2M}{r}} \left[(f_r \bar{p} + p \bar{f}_r) \partial_r \left(1 - \frac{2M}{r} \right) + \left(1 - \frac{2M}{r} \right) (f_{rr} \bar{p} + p \bar{f}_{rr} + f_r \bar{p}_r + p_r \bar{f}_r) \right]
 \end{aligned}$$

$$\begin{aligned}
 &= \bar{p} \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[f_r V_r e^V \sqrt{1 - \frac{2M}{r}} + f_{rr} e^V \sqrt{1 - \frac{2M}{r}} + \frac{2f_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] \\
 &\quad + p \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[\bar{f}_r V_r e^V \sqrt{1 - \frac{2M}{r}} + \bar{f}_{rr} e^V \sqrt{1 - \frac{2M}{r}} + \frac{2\bar{f}_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] \\
 &\quad + \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[V_r e^V \sqrt{1 - \frac{2M}{r}} (f_r \bar{p} + p \bar{f}_r) \right] - \frac{2M_t \mu_0}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \\
 &\quad + \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[e^V (f_r \bar{p} + p \bar{f}_r) \left(1 - \frac{2M}{r}\right)^{-1/2} \partial_r \left(1 - \frac{2M}{r}\right) + e^V \sqrt{1 - \frac{2M}{r}} (f_r \bar{p}_r + p_r \bar{f}_r) \right] \\
 &= \bar{p} \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[f_r V_r e^V \sqrt{1 - \frac{2M}{r}} + f_{rr} e^V \sqrt{1 - \frac{2M}{r}} + \frac{2f_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] \\
 &\quad + p \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[\bar{f}_r V_r e^V \sqrt{1 - \frac{2M}{r}} + \bar{f}_{rr} e^V \sqrt{1 - \frac{2M}{r}} + \frac{2\bar{f}_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] \\
 &\quad + \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[V_r e^V \sqrt{1 - \frac{2M}{r}} (f_r \bar{p} + p \bar{f}_r) \right] - \frac{2M_t \mu_0}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \\
 &\quad + \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[2e^V (f_r \bar{p} + p \bar{f}_r) \partial_r \left(\sqrt{1 - \frac{2M}{r}} \right) + e^V \sqrt{1 - \frac{2M}{r}} (f_r \bar{p}_r + p_r \bar{f}_r) \right] \\
 \mu_t &= \bar{p} \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[\partial_r \left(f_r e^V \sqrt{1 - \frac{2M}{r}} \right) + \frac{2f_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] \\
 &\quad + p \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[\partial_r \left(\bar{f}_r e^V \sqrt{1 - \frac{2M}{r}} \right) + \frac{2\bar{f}_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] - \frac{2M_t \mu_0}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \\
 &\quad + \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[\bar{f}_r \partial_r \left(p e^V \sqrt{1 - \frac{2M}{r}} \right) + f_r \partial_r \left(\bar{p} e^V \sqrt{1 - \frac{2M}{r}} \right) \right] \tag{118b}
 \end{aligned}$$

Setting (118a) equal to (118b), we obtain

$$\begin{aligned}
 &\mu_0 \left[\bar{p} \left(f e^V \sqrt{1 - \frac{2M}{r}} + \frac{p_t}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \right) + p \left(\bar{f} e^V \sqrt{1 - \frac{2M}{r}} + \frac{\bar{p}_t}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \right) \right. \\
 &\quad \left. - \frac{2M_t}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) + \left(1 - \frac{2M}{r}\right) \frac{\bar{f}_r \partial_r \left(p e^V \sqrt{1 - \frac{2M}{r}} \right) + f_r \partial_r \left(\bar{p} e^V \sqrt{1 - \frac{2M}{r}} \right)}{\Upsilon^2} \right] \\
 &= \bar{p} \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[\partial_r \left(f_r e^V \sqrt{1 - \frac{2M}{r}} \right) + \frac{2f_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] \\
 &\quad + p \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[\partial_r \left(\bar{f}_r e^V \sqrt{1 - \frac{2M}{r}} \right) + \frac{2\bar{f}_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] - \frac{2M_t \mu_0}{r} \left(\frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) \\
 &\quad + \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left[\bar{f}_r \partial_r \left(p e^V \sqrt{1 - \frac{2M}{r}} \right) + f_r \partial_r \left(\bar{p} e^V \sqrt{1 - \frac{2M}{r}} \right) \right]
 \end{aligned}$$

$$\begin{aligned}
 & \bar{p} \left(\Upsilon^2 f e^V \left(1 - \frac{2M}{r} \right)^{-1/2} + p_t \right) + p \left(\Upsilon^2 \bar{f} e^V \left(1 - \frac{2M}{r} \right)^{-1/2} + \bar{p}_t \right) \\
 &= \bar{p} \left[\partial_r \left(f_r e^V \sqrt{1 - \frac{2M}{r}} \right) + \frac{2f_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] + p \left[\partial_r \left(\bar{f}_r e^V \sqrt{1 - \frac{2M}{r}} \right) + \frac{2\bar{f}_r e^V}{r} \sqrt{1 - \frac{2M}{r}} \right] \\
 \\
 0 &= \bar{p} \left[-p_t + e^V \left(-\Upsilon^2 f \left(1 - \frac{2M}{r} \right)^{-1/2} + \frac{2f_r}{r} \sqrt{1 - \frac{2M}{r}} \right) + \partial_r \left(f_r e^V \sqrt{1 - \frac{2M}{r}} \right) \right] \\
 &+ p \left[-\bar{p}_t + e^V \left(-\Upsilon^2 \bar{f} \left(1 - \frac{2M}{r} \right)^{-1/2} + \frac{2\bar{f}_r}{r} \sqrt{1 - \frac{2M}{r}} \right) + \partial_r \left(\bar{f}_r e^V \sqrt{1 - \frac{2M}{r}} \right) \right]
 \end{aligned} \tag{118c}$$

Note that, since V and M are real valued, the second term above is the complex conjugate of the first term. Then by (54) equation (118c) holds.

Next, we consider (104b). We first differentiate (117b) with respect to t and then compare it to (104b) after substituting (117), (50), and (51). This yields,

$$\begin{aligned}
 \rho_t &= \partial_t \left[\frac{2\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \text{Re}(f_r \bar{p}) \right] \\
 &= \frac{\mu_0}{\Upsilon^2} \left[-\frac{4M_t}{r} \text{Re}(f_r \bar{p}) + \left(1 - \frac{2M}{r} \right) (f_{rt} \bar{p} + \bar{f}_{rt} p + f_r \bar{p}_t + \bar{f}_r p_t) \right] \\
 &= \frac{\mu_0}{\Upsilon^2} \left\{ -\frac{4M_t}{r} \text{Re}(f_r \bar{p}) + \left(1 - \frac{2M}{r} \right) \left[\bar{p} \partial_r \left(p e^V \sqrt{1 - \frac{2M}{r}} \right) \right. \right. \\
 &\quad \left. \left. + p \partial_r \left(\bar{p} e^V \sqrt{1 - \frac{2M}{r}} \right) + f_r \bar{p}_t + \bar{f}_r p_t \right] \right\} \\
 &= \frac{\mu_0}{\Upsilon^2} \left\{ -\frac{4M_t}{r} \text{Re}(f_r \bar{p}) + \left(1 - \frac{2M}{r} \right) \left[\bar{p} \partial_r \left(p e^V \sqrt{1 - \frac{2M}{r}} \right) \right. \right. \\
 &\quad \left. \left. + \bar{p}_r p e^V \sqrt{1 - \frac{2M}{r}} + |p|^2 \partial_r \left(e^V \sqrt{1 - \frac{2M}{r}} \right) + f_r \bar{p}_t + \bar{f}_r p_t \right] \right\} \\
 &= \frac{\mu_0}{\Upsilon^2} \left\{ -\frac{4M_t}{r} \text{Re}(f_r \bar{p}) + \left(1 - \frac{2M}{r} \right) \left[\partial_r \left(|p|^2 e^V \sqrt{1 - \frac{2M}{r}} \right) \right. \right. \\
 &\quad \left. \left. + |p|^2 \partial_r \left(e^V \sqrt{1 - \frac{2M}{r}} \right) + f_r \bar{p}_t + \bar{f}_r p_t \right] \right\}
 \end{aligned} \tag{119a}$$

From (104b), we obtain

$$\begin{aligned}
 \rho_t &= e^V \sqrt{1 - \frac{2M}{r}} \left(V_r(\mu + P) + P_r + \frac{2(P - Q)}{r} \right) - \frac{2\rho M_t}{r} \left(1 - \frac{2M}{r} \right)^{-1} \\
 &= e^V \sqrt{1 - \frac{2M}{r}} \left[\frac{2\mu_0 V_r}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) (|f_r|^2 + |p|^2) \right. \\
 &\quad \left. + \mu_0 \partial_r \left(-|f|^2 + \left(1 - \frac{2M}{r} \right) \frac{|f_r|^2 + |p|^2}{\Upsilon^2} \right) + \frac{4\mu_0 |f_r|^2}{r\Upsilon^2} \left(1 - \frac{2M}{r} \right) \right] - \frac{4\mu_0 M_t}{r\Upsilon^2} \text{Re}(f_r \bar{p}) \\
 &= \frac{\mu_0 e^V}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \left\{ 2V_r |f_r|^2 \sqrt{1 - \frac{2M}{r}} + 2V_r |p|^2 \sqrt{1 - \frac{2M}{r}} - \Upsilon^2 (f_r \bar{f} + \bar{f}_r f) \left(1 - \frac{2M}{r} \right)^{-1/2} \right. \\
 &\quad \left. + \left(1 - \frac{2M}{r} \right)^{-1/2} \partial_r \left(1 - \frac{2M}{r} \right) (|f_r|^2 + |p|^2) + \sqrt{1 - \frac{2M}{r}} \partial_r (|f_r|^2 + |p|^2) \right. \\
 &\quad \left. + \frac{4|f_r|^2}{r} \sqrt{1 - \frac{2M}{r}} \right\} - \frac{4\mu_0 M_t}{r\Upsilon^2} \text{Re}(f_r \bar{p}) \\
 &= \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \left\{ \bar{f}_r \left[f_r V_r e^V \sqrt{1 - \frac{2M}{r}} - e^V \Upsilon^2 f \left(1 - \frac{2M}{r} \right)^{-1/2} + \frac{2e^V f_r}{r} \sqrt{1 - \frac{2M}{r}} \right] \right. \\
 &\quad \left. + f_r \left[\bar{f}_r V_r e^V \sqrt{1 - \frac{2M}{r}} - e^V \Upsilon^2 \bar{f} \left(1 - \frac{2M}{r} \right)^{-1/2} + \frac{2e^V \bar{f}_r}{r} \sqrt{1 - \frac{2M}{r}} \right] \right. \\
 &\quad \left. + 2V_r e^V |p|^2 \sqrt{1 - \frac{2M}{r}} + 2|p|^2 e^V \partial_r \left(\sqrt{1 - \frac{2M}{r}} \right) + e^V \sqrt{1 - \frac{2M}{r}} \partial_r (|p|^2) \right. \\
 &\quad \left. + 2|f_r|^2 e^V \partial_r \left(\sqrt{1 - \frac{2M}{r}} \right) + e^V \sqrt{1 - \frac{2M}{r}} (f_{rr} \bar{f}_r + f_r \bar{f}_{rr}) \right\} - \frac{4\mu_0 M_t}{r\Upsilon^2} \text{Re}(f_r \bar{p}) \\
 &= \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r} \right) \left\{ \bar{f}_r \left[\partial_r \left(f_r e^V \sqrt{1 - \frac{2M}{r}} \right) + e^V \left(-\Upsilon^2 f \left(1 - \frac{2M}{r} \right)^{-1/2} + \frac{2f_r}{r} \sqrt{1 - \frac{2M}{r}} \right) \right] \right. \\
 &\quad \left. + f_r \left[\partial_r \left(\bar{f}_r e^V \sqrt{1 - \frac{2M}{r}} \right) + e^V \left(-\Upsilon^2 \bar{f} \left(1 - \frac{2M}{r} \right)^{-1/2} + \frac{2\bar{f}_r}{r} \sqrt{1 - \frac{2M}{r}} \right) \right] \right. \\
 &\quad \left. + |p|^2 \partial_r \left(e^V \sqrt{1 - \frac{2M}{r}} \right) + \partial_r \left(|p|^2 e^V \sqrt{1 - \frac{2M}{r}} \right) \right\} - \frac{4\mu_0 M_t}{r\Upsilon^2} \text{Re}(f_r \bar{p}) \tag{119b}
 \end{aligned}$$

Equating (119a) and (119b) yields

$$\begin{aligned}
 & \frac{\mu_0}{\Upsilon^2} \left\{ -\frac{4M_t}{r} \operatorname{Re}(f_r \bar{p}) + \left(1 - \frac{2M}{r}\right) \left[\partial_r \left(|p|^2 e^V \sqrt{1 - \frac{2M}{r}} \right) + |p|^2 \partial_r \left(e^V \sqrt{1 - \frac{2M}{r}} \right) + f_r \bar{p}_t + \bar{f}_r p_t \right] \right\} \\
 &= \frac{\mu_0}{\Upsilon^2} \left(1 - \frac{2M}{r}\right) \left\{ \bar{f}_r \left[\partial_r \left(f_r e^V \sqrt{1 - \frac{2M}{r}} \right) + e^V \left(-\Upsilon^2 f \left(1 - \frac{2M}{r}\right)^{-1/2} + \frac{2f_r}{r} \sqrt{1 - \frac{2M}{r}} \right) \right] \right. \\
 &\quad \left. + f_r \left[\partial_r \left(\bar{f}_r e^V \sqrt{1 - \frac{2M}{r}} \right) + e^V \left(-\Upsilon^2 \bar{f} \left(1 - \frac{2M}{r}\right)^{-1/2} + \frac{2\bar{f}_r}{r} \sqrt{1 - \frac{2M}{r}} \right) \right] \right. \\
 &\quad \left. + |p|^2 \partial_r \left(e^V \sqrt{1 - \frac{2M}{r}} \right) + \partial_r \left(|p|^2 e^V \sqrt{1 - \frac{2M}{r}} \right) \right\} - \frac{4\mu_0 M_t}{r \Upsilon^2} \operatorname{Re}(f_r \bar{p}) \\
 \\
 &0 = \bar{f}_r \left[-p_t + e^V \left(-\Upsilon^2 f \left(1 - \frac{2M}{r}\right)^{-1/2} + \frac{2f_r}{r} \sqrt{1 - \frac{2M}{r}} \right) + \partial_r \left(f_r e^V \sqrt{1 - \frac{2M}{r}} \right) \right] \\
 &\quad + f_r \left[-\bar{p}_t + e^V \left(-\Upsilon^2 \bar{f} \left(1 - \frac{2M}{r}\right)^{-1/2} + \frac{2\bar{f}_r}{r} \sqrt{1 - \frac{2M}{r}} \right) + \partial_r \left(\bar{f}_r e^V \sqrt{1 - \frac{2M}{r}} \right) \right] \\
 &\hspace{25em} (119c)
 \end{aligned}$$

As before, since M and V are real valued, the second term above is the complex conjugate of the first term. Then by equation (54), equation (119c) holds. Then equations (54) and (58) imply equation (104) holds, which completes the proof. $\square$

References

- [1] Éricourgoulhon, 3 + 1 *Formalism in General Relativity: Bases of Numerical Relativity*, Springer, Berlin, 2012.
- [2] Miguel Alcubierre, *Introduction to 3 + 1 Numerical Relativity*, Oxford University Press, Oxford, 2008.
- [3] Miguel Alcubierre, Gabrielle Allen, Bernd Brügmann, Edward Seidel, and Wai-Mo Suen, *Towards an Understanding of the Stability Properties of the 3 + 1 Evolution Equations in General Relativity*, Phys. Rev. D **62** (2000Nov), 124011.
- [4] Richard Arnowitt, Stanley Deser, and Charles Misner, *Republication of: The Dynamics of General Relativity*, General Relativity and Gravitation **40** (2008), 1997–2027. 10.1007/s10714-008-0661-1.
- [5] Thomas W. Baumgarte and Stuart L. Shapiro, *Numerical Relativity: Solving Einstein's Equations on the Computer*, Cambridge University Press, 2010.
- [6] C. Bona, T. Ledvinka, and C. Palenzuela, *3+1 Covariant Suite of Numerical Relativity Evolution Systems*, Phys. Rev. D **66** (2002Oct), no. 8, 084013.
- [7] C. Bona and J. Massó, *Hyperbolic Evolution System for Numerical Relativity*, Phys. Rev. Lett. **68** (1992Feb), 1097–1099.
- [8] Hubert Bray, Sean Hayward, Marc Mars, and Walter Simon, *Generalized Inverse Mean Curvature Flows in Spacetime*, Commun. Math. Phys. **272** (2007), 119–138.
- [9] Hubert L. Bray, *On Dark Matter, Spiral Galaxies, and the Axioms of General Relativity*, [arXiv:submit/0028100 [astro-ph.GA]] (2010).

- [10] Matthew W. Choptuik, *UT Austin PHY387N Lecture Notes*, 1998.
- [11] Isabel Cordero-Carrion, Jose Maria Ibenez, and Juan Antonio Morales-Lladosa, *Maximal slicings in Spherical Symmetry: Local Existence and Construction*, J. Math. Phys. **52** (2011), no. 112501.
- [12] S.W. Hawking, *Gravitational Radiation in an Expanding Universe*, J. Math. Phys. **9** (1968), 598–604.
- [13] Sergey Mashchenko, Alison Sills, and H. M. Couchman, *Constraining Global Properties of the Draco Dwarf Spheroidal Galaxy*, The Astrophysical Journal **640** (2006), no. 1, 252.
- [14] Robert M. Wald, *General Relativity*, The Univeristy of Chicago Press, Chicago, 1984.